On the γ_5 -Gauge Invariance and Helicity Conservation in High-Energy Meson-Baryon Scattering.

M. SEWERYŃSKI

*International Atomic Energy Agency
United Nations Educational Scientific and Cultural Organization
International Centre for Theoretical Physics - Miramare (Trieste)*

(*Lett. Nuovo Cimento*, **4**, 819 (1970))

Page 820, footnote (**),

please replace $\frac{d\sigma(\lambda_3, \lambda_1)}{d\Omega}$ by $\frac{d\sigma(W, \theta)}{d\Omega}$ everywhere;

page 822, lines 10 and 11 should read:

$$\begin{aligned} \phi_2 \pm \phi_3 = & -\frac{1}{M\sqrt{6}} \left[q' W \left(G_1^- \cos \frac{\theta}{2} \pm G_2^- \sin \frac{\theta}{2} \right) + \right. \\ & \left. + (q' \omega_q + qE' \cos \theta) \left(G_1^+ \cos \frac{\theta}{2} \pm G_2^+ \sin \frac{\theta}{2} \right) \mp \frac{M}{2} q \left(G_1^+ \cos \frac{\theta}{2} \mp G_2^+ \sin \frac{\theta}{2} \right) \right]; \end{aligned}$$

page 823, line 2 from below should read:

$$\eta_1^{+\frac{1}{2}} = -\eta_2^{-\frac{1}{2}} = -\frac{1}{M\sqrt{6}} [q'(\omega_q + \omega_{q'}) + E'(q' + q \cos \theta)];$$

page 824, line 2 should read:

$$\xi_1^{+\frac{1}{2}} = \xi_2^{-\frac{1}{2}} = +\frac{1}{M\sqrt{6}} [q'(\omega_q - \omega_{q'}) + E'(q' - q \cos \theta)].$$