The Pion Trajectory in the Processes $\pi^+\text{p} \rightarrow \rho^0\Delta^{++}$ and $\text{K}^+\text{p} \rightarrow \text{K}_{890}^{*0}\Delta^{++}$.

M. MARKYTAN and P. SCHMID

Institut für Hochenergiephysik der Österreichischen Akademie der Wissenschaften - Wien

(*Lett. Nuovo Cimento*, **3**, 51 (1970))

Page 53, fifth line from below: $\beta(\pi) = 24.0$, $\beta(\text{K}) = 20.4$ instead of $\beta(\pi) = 12.0$, $\beta(\text{K}) = 10.2$.

Additional fits have been done with $|t - (m_\Delta - m_N)^2|$ replaced by $|t - (m_\Delta - m_N)^2|^\kappa$ in eq. (5). The results are for $\kappa = 0$ (-1):

$$R_\rho = 2.36 \text{ (2.40)}, \quad R_{\text{K}^*} = 1.14 \text{ (1.10)}, \quad R_\Delta = 1.18 \text{ (1.18)} \text{ (GeV/c)}^{-1},$$

$$\alpha' = 0.54 \text{ (0.49)} \text{ (GeV/c)}^{-2}, \quad s_0 = 0.2 \text{ (0.17)} \text{ (GeV)}^2,$$

$$\beta(\pi) = 43.4 \text{ (63.8)}, \quad \beta(\text{K}) = 38.0 \text{ (52.8)}, \quad \chi^2 = 115.0 \text{ (117.6)}$$

for 50 degrees of freedom.