Addendum

Article 'Streptococcal pharyngitis : clinical clues in diagnosis' by H. Paramesh, M.D. 51:177-179, 1984.

Page 177 : Read 'If the culture was positive for group 'A' beta hemolytic Streptococci, benzathine penicillin 600,000 units was given I.M. to children less than 5 years and 1.2 million units to children above 5 years' *instead of* If the culture was positive for group 'A' beta hemolytic Streptococci, benzathine penicillin 6000,000 units was given I.M. to children less than 5 years and 1.2 million units to children above 5 years'.

Page 178 : Read In place where there are no facilities to culture, on clinical grounds only, on may administer single does of intramuscular benzathine penicillin G 600,-

000 units in children less than 5 years and 1,200,000 units over 5 years' *instead of* In place where there are no facilities to culture, on clinical grounds only, one may administer single dose of intramuscular benzethine penicillin G 600,000 units in children less than 5 years and 1 m 20000 units over 5 years'.

Article 'D-penicillamine in the treatment of Indian childhood cirrhosis-a preliminary report' by B.S. Tomar et al published in Indian Journal of Pediatrics, 50:613-618, 1983 table 6 was inadvertently left out. It is being reproduced now. (The error is regreted-Editor).

Table VI. Hepatic copper content in ICC patients receiving D-penciltamine (group B)

Grades of ICC	Hepatic copper content (ug/g dry matter)														
	At the time of starting of treatment			On 3rd month of follow up			On 6th month of follow up			On 9th month of follow up			On 12th month of follow up		
	Range	Mean	S.D.	Range	Mean	S.D.	Range	Mean	S.D.	Range	Mean	S.D.	Range	Mean	S.D.
I	80-100	85.7	± 7.67	82-102	87.75	± 8.03	76-96	81.55	± 7.39	68-82	71.25	± 5.45	62-74	70.2	± 4.74
II	470-510	491.64	± 11.54	470-502	486.56	± 8.01	470-500	479.92	± 6.23	422-480	433.04	± 25.13	408-440	422.64	± 8.23
III	920-1200	974.8	± 74.94	918-1202	986	± 82.86	916-1200	983.32	± 84.49	910-1008	954.55	± 95.94	908-1002	946.8	± 33.38