- Muscle biopsy and muscle culture in a patient with gout. *J. Neurol. Sci.* 47: 191-202; 1980.
18. Sammons, D. W.; Chilson, O. P. AMP deaminase: stage-specific isozymes in differentiating chick muscle. *Arch. Biochem. Biophys.* 191: 561-570; 1978.
 19. Meienhofer, M. C. Askanas, V.; Proux-Daegelen, M. S.; Dreyfus, J. C.; Engel, W. K. Muscle-type phosphorylase activity present in muscle cells cultured from three patients with myophosphorylase deficiency. *Arch. Neurol.* 34: 779-781; 1977.
 20. Sato, K.; Imai, F.; Hatayama, I.; Roelofs, R. I. Characterization of glycogen phosphorylase isoenzymes present in cultured skeletal muscle from patients with McArdle's disease. *Biochem. Biophys. Res. Commun.* 78: 663-668; 1977.
 21. DiMauro, S.; Arnold, S.; Miranda, A.; Rowland, L. P. McArdle disease: the mystery of reappearing phosphorylase activity in muscle culture—a fetal isoenzyme. *Ann. Neurol.* 3: 60-66; 1978.
 22. Osame, M.; Engel, A. G.; Rebouche, C. J.; Scott, R. E. Freeze-fracture electron microscopic analysis of plasma membranes of cultured muscle cells in Duchenne dystrophy. *Neurology (NY)* 31: 972-979; 1981.

This study was supported by research grants AM27451 and NS06277 from the National Institutes of Health and by a Research Center Grant from the Muscular Dystrophy Association.

ERRATA

Amino Acid and Energy Requirements for Rat Hepatocytes in Primary Culture.

Per E. Schwarze, Anne E. Solheim, and Per O. Seglen

Vol. 18, No. 1, page 44. In Table 1 the legend for Final Concentrations should be μM , not mM ; the Final Concentration for KCl under μM should be 5000.00, not 500.00. On page 45, *Chemicals.* should be [^{14}C]Valine (CFB 75; 50 $\mu Ci/ml$), not [^{14}C]Valine (CFB 75; 50 $\mu Ci/mol$).