**Condition pour la fonction de convergence
dans un modèle composé de particules élémentaires.**

C. BOURRELY et ROAELINA ANDRIAMBOLOLONA

Centre de Physique Théorique, CNRS - Marseille

(Nuovo Cimento, 58 A, 205 (1968))

A la place du titre lire :

Conditions sur la fonction de convergence dans un modèle composé de particules élémentaires.

Absorption Corrections for the Double Peripheral Model.

J. L. SCHONFELDER

Physics Department, Imperial College - London

(Nuovo Cimento, 58 A, 221 (1968))

The angles $(\theta_3 - \theta'_3)$ etc., appearing on page 228 and especially in eq. (17) should be replaced by $\cos \alpha_3$ etc., where $\cos \alpha_3$ is given by

$$\cos \alpha_3 = \cos \theta'_3 \cos \theta_3 + \sin \theta_3 \sin \theta'_3 \cos \varphi'_3$$

and similarly for $\cos \alpha_1$ and $\cos \alpha_2$.

Thus eq. (17) should read

$$\begin{aligned} \langle \lambda_1 \lambda_2 \lambda_3 | T | \lambda_a \lambda_b \rangle &= \langle \lambda_1 \lambda_2 \lambda_3 | V | \lambda_a \lambda_b \rangle + \\ &+ \frac{p_a A_{(ab)}}{16\pi W} \int d \cos \theta' \langle \lambda_1 \lambda_2 \lambda_3 | V(S_{ab}, S_{12}, S_{23}, \cos(\Theta_3 - \theta'), \varphi^3) | \lambda_a \lambda_b \rangle \exp[b_{(ab)} p_a^2 (\cos \theta' - 1)] + \\ &+ \frac{p_1^{(3)} A_{(12)}}{8W_3(2\pi)^2} \int d \cos \theta'_3 d \varphi'_3 \exp[b_{(12)} p_1^{(3)2} (\cos \alpha_3 - 1)] \langle \lambda_1 \lambda_2 \lambda_3 | V(S_{1b}, S_{12}, \cos \Theta_3, \cos \theta'_3, \varphi'_3) | \lambda_a \lambda_b \rangle + \end{aligned}$$

+ two similar terms with indices 123 cyclicly interchanged.