SUMMARY

A case of pemphigus neonatorum treated with cortisone has been reported. There was no response to ilotycin and terramycin therapy. The response to cortisone therapy was dramatic. The possible mode of action of the hormone in pemphigus has been discussed.

REFERENCES :

1. BRENNAN, J. C. AND MONTGOMERY, H.—Pemphigus and other bullous dermatoses, correlation of clinical and pathological findings, *J.A. Derm.*, **21** : 349-91, 1953.
2. CANIZARES, O.—Pemphigus, *Arch. Derm. Syph.*, **65** : 499, Chicago.
3. COSTELLO, M. J.—*Arch. Derm. Syph.*, **65** : 498, Chicago.
4. FISHER, ISADOR.—Pemphigus Vulgaris, *A.M.A. Arch. Derm.*, **74** : 50-58, 1956.
5. KIERLAND, R. R., O'LEARY, P. A., BRUNSTING, L. A. AND DIDCOCK, J. W.—Cortisone and Corticotrophin in Dermatology, *J.A.M.A.*, **148** : 23, 1952.
6. LEVER, W. L.—Pemphigus, *Medicine*, **32** : 1, 1953.

ERRATA

Indian J. Pediat. **24** : 151, 1957 Ref. 3, should read: BERT, J.M.—Quoted by WHITESIDE AND HOLMELEY, and p. 150, line 26 should read SHELLING.^{15a} Insert below Ref. 15, page 151 the following: 15a. *Idem* quoted by FELDMAN AND DAVENPORT.