a universality also for M_1 . In fact we find $g_{M_1 BB}/g_{M_1 PP} = \frac{3}{2}$ (notice that ξ cancels out in this ratio). Even though M_1 need not be a particle, this fixes the universality of the P-trajectory and we get $P_N/P_\pi = \frac{3}{2}$ in eq. (2). It is not clear what the assumption that M_{144} and never $\text{Tr}(M_{144})$ should appear means at a physical level. It is to be considered for the time being as a postulate of the Okubo type that enables us to derive the ratio and the justification for which is probably to be found within a scheme transcending $U_6 \times U_6$ much like the justification for Okubo's prescription was found beyond SU_3 .

We still wish to make some remarks about the Pomeranchukon. Since it is in the representation (1, 1) of $U_6 \times U_6$, it is not to be thought of as a bound system in the $9\bar{9}$ channel (the same is true for the vacuum itself). This may well account for its very small slope compared to the $R_{\pm t}$ trajectories as required by fits to diffraction scattering experiments^(*). For all we can tell there is no reason why particles should at all lie on the P-trajectory. We can paraphrase CHKN by concluding that since we assume the P-trajectory to belong to a $U_6 \times U_6$ singlet, we find it natural that its properties should be quantitatively and qualitatively different from those of other poles of even signature.

(*) See e.g. R. PHILLIPS and W. RARITA: *Phys. Rev.*, **139**, B 1336 (1965).

Experimental Tests of SU_{6W} in Nucleon-Antinucleon Annihilation at Rest.

A. BETTINI

Istituto di Fisica « Galileo Galilei » dell'Università - Padova
Istituto Nazionale di Fisica Nucleare - Sezione di Padova

(*Nuovo Cimento*, **44 A**, 285 (1966))

In Table I:

<i>errata</i>		<i>corrigé</i>	
$^1S_0 \text{ p}\bar{\text{p}} \rightarrow \bar{\text{K}}^{*0} \text{K}^0$	8	$^1S_0 \text{ p}\bar{\text{p}} \rightarrow \bar{\text{K}}^{*0} \text{K}^0$	0.5
$\bar{\text{K}}^{*-} \text{K}^+$	0.5,	$\text{K}^{*-} \text{K}^+$	8.