which should be compared with the more recent experimental determination ⁽⁹⁾

$$(8) \quad \tau_{\text{exp.}} \cong 4 \cdot 10^{-16} \text{ s.}$$

Thus we see that the agreement of the theory with experiment has improved and this may be considered an argument in favour of the symmetrical inter-

action (2), at least in what refers to nucleons and cascade particles. The question of the sign of the terms in Σ and Λ (and thus the choice between the two theories) should be decided from other experimental results, in particular, from the scattering of π by protons.

* * *

We should express our acknowledgements to the U.S. Office of Naval Research who made possible our attendance to the Rochester Conference.

⁽⁹⁾ J. OREAR as referred by L. W. ALVAREZ in the Rochester Conference (April 1957) and private communication.

EMENDATION

R. WEINER — **Nuclear Isomeric Shift on Spectral Lines**, *Nuovo Cimento*, **4**, 1587 (1956).

Some mistakes are contained in the above paper and they should be emended as follows:

page 1587 4th line from the bottom	= e	should read	= $-e$
page 1588 14th line	= e	»	» = $-e$
page 1588 17th line	the last factor r^{2n_r+l+1}	»	» r^{2n_r+2l+1}
page 1589 2nd and 3rd lines	... state;	»	» ... state).
		and delete the rest of the paragraph.	
page 1589 12th line	≈ -5	should read	$\approx +5$