ERRATA

Primary Aldosteronism in Pregnancy – Should it be Treated Surgically?

E. Aboud, M. De Swiet, H. Gordon

The following Table was omitted when published in the October issue

TABLE I
SUMMARY OF ALL CASES PUBLISHED SINCE 1966

Publications	No. of cases	Improved Preg.	Deteriorated Preg.	Postpartum outcome.	Comment
Gordon et al 1967	1	–	–	–	Adenoma removed at 18 weeks
Biglieri and Slaton 1967	1	1	–	#	–
Levy and Marx 1971	1	–	1	**	–
Aoi et al 1978	2	2	–	Both worse	–
Hammond et al 1982	1	–	1	**	–
Elterman et al 1983	1	–	1	Worse	–
Merrill et al 1984	1	–	1	Worse	–
Colton et al 1984	1	–	–	–	Termination of pregnancy first trimester
Lotgering et al 1986	1	–	1	Worse	–
Neerhof et al 1991	1	–	1	**	–
TOTAL	11	3	6	5	2

Initially deteriorated then improved

** No information provided

Recurrent Laryngeal Nerve Protection in Thyroidectomy

D. S. O’Riordan, M. P. Brady

Ir. J. Med. Sci. 1995; 164: 265-266

Paragraph 2, Line 18. The sentence beginning “Inferior thyroid veins” should read:
“Inferior thyroid veins should *not* be ligated early in the procedure as an occasional anteriorly placed RLN may be endangered”.

Comparison of Two Fluorescent Probes for the Measurement of Erythrocyte Membrane Fluidity in Renal Dialysis Patients

E. McClean, L. T. McGrath, G. P. R. Archbold

Ir. J. Med. Sci. 1995; 164: 289-292

The formula for calculating the fluorescence polarisation (p) should read:

$$p = \frac{I_{\text{par}} - GI_{\text{per}}}{I_{\text{par}} + GI_{\text{per}}}$$

I_{par} = fluorescence parallel to the plane of polarisation

I_{per} = fluorescence perpendicular to the plane of polarisation