LETTERS TO THE EDITOR

Victoria Hospital,
London, Ontario,
December 20, 1963

SIR

I read with interest the comprehensive paper by Drs Snow and Vandewater, entitled "Gall-Stones and Anaesthetics"* The relationship of gall bladder disease to heart disease is clearly elucidated, but perhaps a few remarks on the implications of anaesthesia in this sort of case are not unwarranted These comments involve mainly the part of the coeliac plexus reflex plays in the aggravation of cardiac disease both preceding and during surgery on the biliary tract Indeed the authors do describe "reflex vagal vasoconstriction of the coronary arteries," abolished by atropine, in their opening paragraph In addition, Ravdin *et al* † have shown that "calculus cholecystitis" can reflexly cause myocardial ischaemia in otherwise healthy patients, or aggravate symptoms in cardiac patients Distention of the common bile duct reflexly lowers coronary blood flow and may result in anginal symptoms Ravdin was even able to reproduce anginal attacks in a patient postoperatively by repeated distention of the common duct

During surgery, traction or manipulation of upper abdominal structures may elicit the coeliac reflex, resulting in sudden hypotension As coronary artery flow depends on mean aortic blood pressure, the possible effects of this reflex occurring during operation in patients with coronary disease should be remembered The authors state "These patients all tolerate operation well, provided that nothing is done to precipitate further myocardial ischaemia" This includes gentle surgical technique, adequate atropine dosage, ‡ and immediate removal of the operative stimulus if hypotension occurs The hypotension is more profound in poor-risk patients and is enhanced by deeper levels of anaesthesia Tachycardia from atropine administration should be avoided as this may increase the cardiac ischaemia

The authors report a 20 per cent incidence of postoperative myocardial infarctions (5 infarcts in 255 patients) All five patients had pre-existing ischaemic heart disease This figure is in keeping with the observation that "morbidity and mortality in those with coronary artery disease is 2 to 3 times higher than in those without involvement" The anaesthetist might be assisted in the possible further reduction of this complication rate by the use of a postoperative intensive care unit and low viscous dextran infusions intravenously

H B KENNELYSIDE, M D

*Vol 10, no 6, pp 555-566 (1963)

†RAVDIN, I S, FITZ-HUGH, T, JR, WOLFERTH, C C, BARBIERI, E A, & RAVDIN, R G Relation of Gallstone Disease to Angina Pectoris Arch Surg 70 333 (1955)

‡SMITH, B H The Nature and Treatment of the Coeliac-Plexus Reflex in Man Lancet 265 223 (1953)

Glaxo-Allenburys (Canada) Ltd.,
52 Bartor Road,
Weston, Ontario,
November 7, 1963

SIR

I think it might be well worth while bringing the following matter to the attention of the Anaesthetists of Canada, through the pages of your journal

Sealed-glass ampoules of water for injection as marketed in Canada contain nothing but apyrogenic sterile water. Rubber-capped, multi-dose vials, on the other hand, in addition to sterile water contain a preservative, which is commonly 1 to 1½ per cent benzyl alcohol. I learned recently that in one Toronto hospital it is common practice to employ water from rubber-capped vials as a diluent for the anaesthetic agent employed in spinal and epidural anaesthesia. In view of the fact that benzyl alcohol has been shown to be capable of producing nerve degeneration, this would appear to be an unsafe practice. While it is true that the degeneration resulting from benzyl alcohol is marked if 10 per cent solutions are used, and is much less when 5 per cent solutions are employed, the possibility of damage from weaker solutions must certainly exist.

Anaesthetists should, therefore, insist that any water to be used in anaesthesia be supplied in sealed-glass, single-dose ampoules to which no preservative has been added.

H. CLARK BALMER, M.D.
Medical Director