fission and separation and reproduction of the two parts.

Among crystals, the closest analogies to the gene are to be found in the nucleo-proteins that are associated with the viruses. They have somewhat the same property of autosynthesis and reproduction. As is now well known, some of the viruses have been crystallized. Viruses resemble genes in that they can be inactivated by Roentgen-rays, and through radiation they can be given new specificities so that offspring of the treated ova will be different from the parent.

It is remarkable to see that within a few years after

the discovery of genes, a man can write 41 pages and quote 274 authorities on the chemistry of these tiny bodies!

ERRATUM

In the case report of the article by Dr. Leonard Cardon entitled "Generalized Pruritus Due to Carcinoma of the Stomach and Cured by Gastrectomy" which appeared in the February issue of this Journal, it was stated that "At the present writing (9-1-42) he weighs over 100 lbs." However, this date was incorrect, and should have been 12-30-42.

Abstracts of Current Literature

FRANKLIN HOLLANDER, Associate Editor in Charge of Abstracts, New York, N. Y.

ALBERT CORNELL, New York, N. Y.
MAURICE FELDMAN, Baltimore, Md.
CHARLES A. FLOOD, New York, N. Y.
M. H. F. FRIEDMAN, Philadelphia, Pa.
J. DUFFY HANCOCK, Louisville, Ky.
THOMAS A. JOHNSON, Philadelphia, Pa.
ALLEN A. JONES, Buffalo, N. Y.
N. W. JONES, Portland, Ore.
HENRY H. LERNER, Boston, Mass.
PHILLIP LEVITSKY, Montreal, Can.

JESSIE LOVE, Louisville, Ky.
FRANZ J. LUST, New York, N. Y.
IRA A. MANVILLE, Portland, Ore.
ARTHUR E. MEYER, Brooklyn, N. Y.
FRANCIS MURPHY, Milwaukee, Wis.
H. NECHELES, Chicago, Ill.
FRANK NEUWELT, Gary, Indiana
SAM OVERSTREET, Louisville, Ky.
J. F. PESSEL, Trenton, N. J.
DAVID J. SANDWEISS, Detroit, Mich.

RUDOLF SCHINDLER, Chicago, Ill.
MICHAEL SHUTKIN, Milwaukee, Wis.
VIRGIL E. SIMPSON, Louisville, Ky.
H. J. SIMS, Denver, Colo.
HENRY TUMEN, Philadelphia, Pa.
ROBERT TURELL, New York, N. Y.
EDGAR WAYBURN, San Francisco, Calif.
JOHN WILLARD, Philadelphia, Pa.
C. WILMER WIRTS, Philadelphia, Pa.

[Microfilm copies of papers may be obtained from Medicofilm Service of the Army Medical Library at 25c for each complete article, not exceeding 25 pages in length—and 10c for each additional 10 pages or fraction thereof. Prepayment is not requested. Remittance may be made with subsequent orders and in such manner as found most convenient. Address—Medicofilm Service, Army Medical Library, Washington, D. C.]

CLINICAL MEDICINE

MOUTH AND ESOPHAGUS

DINSMORE, R. S.: *Diverticulum of the Esophagus. Cleveland Clin. Quart.*, 9:187, Oct., 1942.

Two cases of esophageal diverticulum occurring in brothers aged 61 years and 51 years are reported. It is not thought this condition is familial. Both of these cases had persisted for four or five years during which time the diverticula had shown evidence of enlargement and symptoms had increased in severity. Both cases were treated with complete success by one stage operations. The author is personally familiar, from his own experience, with single stage and two stage surgical procedures and believes each has its special advantages in selected cases. He prefers the one stage operation as technically easier and safer whenever it can be employed. His technique is carefully described. The report constitutes a very interesting and informative contribution, even to the physician not primarily interested in surgery.—Sam A. Overstreet.

HENRY, T. C.: *Ultero-Membranous Stomatitis. Brit. Med. J.*, 4261:273, Sept., 1942.

Among 90 cases of ultero-membranous stomatitis, 87 were found to be Vincent's disorder and 3 were atypical. Most of the cases of atypical ultero-membranous stomatitis which have been reported bear a superficial resemblance to Vincent's angina but differ from this in the degree of toxemia they cause, the presence of complications and the relatively high mortality. In some instances the cases

of ultero-membranous stomatitis seen represent the oral manifestations of such diseases as memphigus, erythema multiforme, or dermatitis herpetiformis.

The author describes 3 cases of atypical ulcerative stomatitis in which marked prostration and toxemia occurred, associated with conjunctivitis. In two cases there were skin lesions. Streptococci were isolated from the mouth lesions of all the cases, being hemolytic in those with skin lesions and non-hemolytic in the other. None of these cases responded to sulfonamide therapy, although all recovered after transfusions.—Henry Tumen.

PENNES, A. E.: *Leiomyosarcoma of the Esophagus. Am. J. Roent. Rad. Therapy*, 48:336, Sept., 1942.

Pennes reports a case of leiomyosarcoma of the esophagus, which produced a roentgen filling defect resembling carcinoma, except that there was an extrinsic mass in the mediastinum connected with the esophageal tumor. The extension of the tumor into the mediastinum is an outstanding feature of the case. Although intra- and extra-luminal tumors have been noted in the stomach, it is a rare occurrence to demonstrate them in the esophagus. Sarcoma of the esophagus is either an ulcerative or polypoid growth. The tumor is frequently largely extra-luminal.—Maurice Feldman.

GRACE, E. J.: *Control of Massive Esophageal Hemorrhage Secondary to Liver Damage (Cirrhosis) by Ligation of the Coronary Vein and Injection of Sodium Morrhuate. Ann. Surg.*, 116:387, Sept., 1942.

The pathogenesis of esophageal varices is first discussed rather completely. Then the author goes on to discuss the evolution of the various techniques for the control of hemorrhage from bleeding esophageal varices: injection of sclerosing solutions, ligation of the coronary vein of the stomach, splenectomy, omentopexy, etc. He then discusses the case of a far advanced cirrhotic with ascites, with poor liver function, and multiple hemorrhages from bleeding varices. Following celiotomy the coronary vein and several other smaller veins close to the former were ligated and sclerosing solution (sodium morrhuate) was injected. The veins of Sappey on the undersurface of the diaphragm were also ligated. The spleen was bound down due to an old perisplenitis and was not removed. A month after the first operation omentopexy was performed. The patient was then given the benefit of modern knowledge in the way of therapy for liver damage and 2 years after the above surgical procedures the man felt fine, had good liver function, had no ascites and worked daily as an elevator operator. He had had no more gross hemorrhages.—Frank Neuwelt.

STOMACH

MCCLURE, C. W. AND JANKELS, I. R.: *The Clinical Application of Gastroscopy. New Eng. J. Med.*, 227:548, Oct., 1942.

Roentgenology and gastroscopy are complementary methods, and are neither competitive nor antagonistic. Their combined use can reduce con-