LETTERS TO THE EDITOR

SIR:

May I draw to your attention an error of omission in the printing of my letter of August 29, 1969, appearing in the November issue of the Canadian Anaesthetists' Society Journal. In the fourth paragraph, second sentence of my original letter it read, "However, I would point out that the incidence of fires and explosions in North America have been unknown for at least twenty years." The letter as printed omits "the incidence of." This gives one the impression that fires and explosions have not occurred in the last twenty years, whereas I was referring to the lack of data on the incidence. Unfortunately, the error will be the focus, if any, of opposing views to the issue that I wish to raise.

STUART L. VANDEWATER, M.D.
Professor of Anaesthesiology
Queen's University
Kingston, Ontario

November 11, 1969

SIR:

You ought to be congratulated for having published in your Journal (vol. 6, no. 6, November 1969) the excellent letter by Dr. Stuart L. Vandewater in which he takes issue with the ban of flammable anaesthetics as a deliberate administrative step by a hospital. For the purposes of this argument it is immaterial whether this action was being concurred in by the professional people in the institution, as indeed it was in this case. The fact remains that an administrative edict of this kind removes from physicians the discretion of using particular agents which, at least in my humble opinion, in this particular case may yet be advantageous under certain clinical circumstances. It is one thing for a physician or a group of physicians to refrain from using certain drugs or techniques; it is quite another matter for the administration to ban them. In the former instance they still would be available when clinical judgment indicates their use, while in the latter this is no longer possible.

It is somewhat incongruous to pick on flammable anaesthetics because a fire or explosion may be caused, when indeed this is an exceedingly rare occurrence – much rarer than the author of the paper implies. In 1950 Foster Kennedy and others made a great impression with their paper exposing the alleged grave dangers of subarachnoid block (*Surg. Gynec. & Obstet.* 91:385, 1950). Then within a few years Beecher and Tood in their "Study of Deaths Associated with Anesthesia and Surgery" implied that muscle relaxants might be prominent in the causation of anaesthetic deaths. As Dr. Vandewater so rightly says, halothane is still under scrutiny as to its hepatotoxic effects and methoxyflurane has been

implicated in the causation of renal dysfunction. The incidence of complications for any anaesthetic agent and technique far exceeds the number of anaesthetic fires and explosions, yet no outright ban has been considered. No doubt certain physicians have been sufficiently impressed by the supposed evidence to eliminate them from their own practice, but have they been submitted to an entirely unjustified wholesale ban? Not even chloroform, which the majority consider indeed to be hepatotoxic, has been banned by administrative edict, although few of us would use it clinically.

At one point in the paper it is urged that "the use of any other non-anaesthetic volatile agents for any purpose be prohibited unless specifically approved by the fire and safety committee and by the administration." In other words, despite the fact that structural safeguards will no longer be provided, exceptions are possible, as long as presumably they do not involve anaesthetists. Do I see in this the long fatherly arm of some influential surgeon who does not wish to forego his cherished skin-prep?

I am looking forward to the logical sequel of all this. In their concern for the physical safety of patients, the hospital administrators undoubtedly will advise in due course that they have banned all electrical and electronic diagnostic and therapeutic equipment from their operating rooms having been unable to control completely the danger of electrocution. The next step then will be a ban of all other potential physical hazards, which presumably must include surgical instruments. This done, complete safety for the patient will have been assured, since there will be no need for operating rooms at all (by far the cheapest solution). Provided now that side-rails are up on beds and that hot water bottles are filled with cold water, no physical harm can possibly come to any patient.

GORDON M. WYANT, F.R.C.P.(C)
Professor of Anaesthesia
University of Saskatchewan
Saskatoon, Sask.

November 14, 1969