saturation reading provide further reassurance of the adequacy of the collateral circulation. In patients who are cold, I have noticed that a "failed" test may subsequently show adequate circulation.

The test is rapid (30 seconds). There is minimal discomfort to patients and it requires no patient cooperation. It does not depend on hyperaemia as indirect evidence of adequate perfusion. After all, it is conceivable that enough collateral flow could percolate through to cause hyperaemia of the hand but leave the tips of the digit(s) ischaemic. If a negative test is defined as adequate collateral circulation, then by definition there are no false negatives. The incidence of false positives would be interesting. My experience indicates only a handful of positive tests (inadequate collateral circulation) in the past year. As a test that looks directly at the endpoint in question, and gives a quantitative index of saturation to the digits during radial artery occlusion, it may prove to be the most useful.

Winston Wong MD FRCPC
Department of Anaesthesia
St Joseph's Health Centre
Toronto, Ont.

REFERENCE

- 1 Vaghadia H, Schechter M, Sheps S, Jenkins L. Evaluation of a postocclusive reactive circulatory hyperaemia (PORCH) test for the assessment of ulnar collateral circulation. *Can J Anaesth* 1988; 35: 591-8.

REPLY

Thank you for the opportunity of replying to the letter by Dr. Wong. The use of pulse oximetry for the assessment of ulnar collateral circulation has already been described.¹ In addition, Kurki et al.² have studied the effects of radial artery cannulation on digital blood pressure (Finapres TM) and oxygen saturation (SaO₂) during and after cannulation. They studied 22 hands with a normal Allen's test and found that SaO₂ was zero in four patients for 2-17 minutes during and after cannulation. In four additional patients the SaO₂ was zero following insertion of a cannula but returned to control within one minute. In those patients where SaO₂ was zero, no blood flow distal to the cannula could be detected by a Doppler ultrasonic velocity meter. The authors felt that despite indications by Allen's test that there was adequate ulnar collateral flow, all non-invasive monitors (Doppler, Finapres TM, and oximeter) placed distal to the site of cannulation can be affected by changes in blood flow for several minutes. These findings suggest that the use of a pulse oximeter as advocated by Dr. Wong and by Nowak et al. would be subject to the same limitations and therefore be unreliable. Digital occlusion of the radial artery may abolish the pulse oximeter reading and yield a high incidence of false positives unless one were prepared to wait from 2 to 17 minutes in order to determine if the SaO₂ readings would be restored.

Dr. Wong suggests that the objective behind testing for ulnar collateral flow is to ensure adequate perfusion to the digits with

FIGURE 1 Result of a PORCH test demonstrating thenar ischaemia with good flow to the tips of the digits. This patient had a normal Allen's test.

FIGURE 2 Release of radial artery compression after PORCH test restores flow to the thenar area in the same hand as shown in Figure 1. This confirms that the radial artery was an end artery to the thenar area in this patient.

manual occlusion of the radial artery. We would add that it is necessary to ensure adequate perfusion of the thenar area as well as the thumb and four digits because ischaemic injury following radial artery cannulation has been reported to involve

the thenar eminence, thumb and index fingers,³ as well as the pads of the third, fourth and fifth digits.⁴ In fact, angiographic studies by Hasse *et al.*⁵ demonstrated that the radial artery is frequently an end artery to the thenar area. Thus, it is possible that a pulse oximeter applied to the tip of a digit may indicate adequate collateral flow to the tip when in fact the thenar area is at risk of ischaemia should radial occlusion be permanent.

During our studies with the PORCH test we observed that some hands with a normal Allen's test exhibited signs of thenar ischaemia with the PORCH test (see Figures 1 and 2). Such patients would probably not have been identified using pulse oximetry as described by Dr. Wong. What appeals to us about the PORCH test is its ability to assess the end-point of interest to anaesthetists – the response of the collateral circulation in the hand to temporary but intense ischaemia. If the collateral circulation is adequate during performance of the PORCH test, it is probably safe to conclude that it will also be adequate in the event of permanent ischaemia. We would emphasize that the attributes of the PORCH test are based on 14 hands with an abnormal collateral circulation. Further assessments will be necessary before these attributes can be stated with precision.

Himat Vaghadia MB FFRCS FRCP
 Martin T. Schechter MD PhD
 Samuel B. Sheps MD FRCP
 Leonard C. Jenkins MD FRCP
 Department of Anaesthesia, Vancouver General Hospital
 and Department of Health Care and Epidemiology
 University of British Columbia
 Room 3200, Faculty of Medicine
 910 West 10th Avenue
 Vancouver, B.C. V5Z 4E3

REFERENCES

- 1 Nowak GS, Moorthy SS, McNiece WL. Use of pulse oximetry for assessment of collateral arterial flow. *Anesthesiology* 1986; 64: 527.
- 2 Kurki TS, Sanford TJ, TySmith N, Dec-Silver D, Head N. Effects of radial artery cannulation on the function of finger blood pressure and pulse oximeter monitors. *Anesthesiology* 1988; 69: 778–82.
- 3 Mangano DT, Hickey RF. Ischemic injury following uncomplicated radial artery catheterization. *Anesth Analg* 1979; 58: 55–7.
- 4 Baker RJ, Chunprapaph B, Nyhus LM. Severe ischemia of the hand following radial artery catheterization. *Surgery* 1976; 80: 449–57.
- 5 Hasse J, Pusterla C, Cloeren S. Angiographische untersuchungen nach dauer kanulierung der arteria radialis. *Schweiz Med Wochenschr* 1971; 101, 1057–61.

Pharmacokinetics of sufentanil

To the Editor:

The recent report by Fyman *et al.*, "Pharmacokinetics of sufentanil in patients undergoing renal transplantation"¹ deserves comment. Several inconsistencies are present in the report. There is disagreement between the pharmaco-

kinetic variables reported in the table, the pharmacokinetic equation and description contained in the legend of the Figure, and the appearance of the Figure itself:

- 1 The legend states that the Figure is a "sufentanil plasma decay curve fitted to a two-compartment model." However, the data in the Figure appear to be mean data (the authors stated that they only fitted individual data), and do not appear "fitted" at all; the data points are merely connected by lines and demonstrate no scatter about a line of best-fit, which should appear as a smooth curve.
- 2 The equation in the legend suggests that the values of α and β (the exponents in the two terms) are 0.621 min^{-1} and 0.025 min^{-1} , respectively. If the values of α and β half-life from the Table are converted to the corresponding exponents by the transformation $0.693/t_{1/2}$, the results are 0.239 min^{-1} and 0.00394 min^{-1} , respectively. This discrepancy may have arisen from taking the arithmetic mean of the half-lives, when the harmonic mean may have been more appropriate.² However, the gross disagreement between the two estimates of β suggests a more serious error. Conversely, if the parameter values for k_{10} , k_{12} , and k_{21} (from the Table) are applied to standard equations³ to calculate the exponents α and β , above, the results are 0.626 min^{-1} and 0.024 min^{-1} respectively. These compare well with the values of the equation in the Figure legend. However, a β value of 0.024, when converted to half-life, yields a value of approximately 30 min, in complete disagreement with the reported mean half-life of 176 min.
- 3 The terminal (β) half-life (stated in the Table) of 176 min, or about 3 hr, does not correspond to the appearance of the concentration versus time plot. The concentration of sufentanil at 120 min is approximately 0.35 to 0.4 $\text{ng} \cdot \text{ml}^{-1}$; at 360 min, 4 hr later, the concentration appears virtually unchanged as do the concentrations at intervening time points. Yet, the half-life stated in the table would predict a concentration of less than 0.2 $\text{ng} \cdot \text{ml}^{-1}$ at 360 min.

The authors concluded that sufentanil disposition in their population of renal transplant patients were not different from that observed in "healthy" patients, offering the data of Bovill *et al.*⁴ as an appropriate comparison. If one compares the plot of mean sufentanil concentration versus time published by Bovill's group with that of Fyman *et al.*¹ striking differences are evident. The scales of the two graphs are very similar, yet the data of Bovill show a clear, linear decrease in concentration with time after 2 hr, corresponding to a half-life of 164 min, while that of Fyman shows little or no decrease in concentration, with a half-life reported to be 176 min. Also, the mean sufentanil concentrations at 6 hr appear