since the classical signs of tachycardia and tearing are by no means constant and in retrospect have not occurred in some patients who, by their own account, have been awake during all, most, or part of the operation. Hence errors of judgement as to the need for deeper anaesthesia and further analgesics do occur and they happen more frequently the less experienced the anaesthetist.

The rationale frequently advanced for the use of this technique is its lack of toxicity. Even if one were to assume complete absence of inherent toxicity in these combinations, an assumption which might be open to challenge, one must still ask the question whether the rather remote chance and low incidence of complications from more potent inhalation agents justify the enormous transgression against a patient who awakens in the middle of an operation, totally paralyzed and unable to communicate his distress. The enormity of the situation is compounded in those cases in which pain sensation is not entirely obtunded. Such a situation comes dangerously close to vivisection. Is it not time now to reassess such techniques, as medical practice comes under increasingly searching scrutiny by the public at large and we can ill afford failures which are easily avoided by discarding techniques which are potentially inadequate?

GORDON M. WYANT,
Department of Anaesthesia,
University of Saskatchewan

CORRECTION

Alphathesin, A New Steroid Anaesthetic Agent

Jean-Paul Dechene

Volume 23. No. 2. March, 1976. Page 163.

The second sentence in the last paragraph on page 163 should read as follows –
“The optimum dosage is in the vicinity of 50 μ l to 60 μ l per kilogram body weight.”