LETTERS TO THE EDITOR

CARCINOID SYNDROME AND ANAESTHETIC MANAGEMENT

SIR,

We have read with great interest the excellent publication about anaesthesia for the carcinoid syndrome by Miller, *et al.*¹ Our experience with five anaesthetics in three patients with carcinoid syndrome are published in the *Acta Anaesthesiologica Belgica*.²

A very important aspect of anaesthetic management which is not infrequently disregarded is the preoperative treatment of dehydration and hypovolaemia.^{2,3} Preoperatively, these patients often have severe diarrhoea as well as hypertension, both of which will promote dehydration and extreme hypovolaemia. As the human body always tries to get a homeostasis, most of the biochemical values seems quite normal, even if there is a marked hypovolaemia.

The clinical findings of hypovolaemia are poor tissue turgor, dry skin and mucosa, slow capillary refill and a marked decrease in the 24-hour urine production.

Our experience is that, when there is a sufficient circulating blood volume, ensured by vigorous preoperative fluid therapy, fluctuations in blood pressure may occur during handling of the tumour, but these do not necessarily lead to disastrous hypotension.

A further clinical impression is that the greatest value has to be attached to adequate analgesia during the operation. We fully agree with Miller, *et al.* that anaesthesia for patients with the carcinoid syndrome may not be as hazardous as previously reported, provided that patients are adequately prepared.¹

Some decrease in blood pressure and a poor systemic circulation can also be expected after the operation. Careful postoperative management and observation of the patient in an intensive care unit, by the anaesthetist is therefore necessary!

Y.H. Khouw M.D.
J.W. Kleine M.D.
Department of Anaesthesiologie
Ziekenhuis "Ziekenzorg"
Enschede, Nederland

REFERENCES

1. MILLER, R., PATEL, A.U., WARNER, R.R.P. and PARNES, I.H. Anaesthesia for the carcinoid syndrome. *Can. Anaesth. Soc. J.* 25: 240-244 (1978).

2. KLEINE, J.W., KHOUW, Y.H., & HEERES, M.G. Management of anaesthesia during surgery for patients with carcinoid syndrome. *Acta Anaesthesiol. Belgica* 28: 165 (1977).
3. KLEINE, J.W., KHOUW, Y.H. & HEERES, M.G. Hypovolaemia and hypotension with carcinoid syndrome. *Anaesthesiology* 49: 55 (1978).
4. MASON, R.A. & STEANE, P.A. Carcinoid syndrome: its relevance to the anaesthetist. *Anaesthesia* 31: 228-242.
5. MASON, R.A. & STEANE, P.A. Anaesthesia for a patient with carcinoid syndrome. *Anaesthesia* 31: 243-246.
6. PATEL, A.U., MILLER, M.D. & WARNER, R.P. Anaesthesia for ligation of the hepatic artery in a patient with carcinoid syndrome. *Anesthesiology* 47: 303-305 (1977).

THE BAIN CIRCUIT

SIR,

Discussion of the Bain Circuit is always interesting, but surely the time has come for reconsideration of low inflow semiclosed carbon dioxide absorption systems. The flow often necessary for the Bain, when combined with a mechanical ventilator, presents environmental control arrangements with an unnecessary and sometime unmanageable burden. Most anaesthetists using narcotic supplemented nitrous oxide techniques consider such flows wise and the Bain system a convenient way of dealing with them. However, much lower inflows can easily be used for a narcotic-nitrous oxide-enflurane or halothane technique. Indeed, available information leads one to question the routine use of nitrous oxide in such a drug combination. Such questions relate to the pharmacological effect of nitrous oxide under the prevailing circumstances, oxygen supply failure, air filled spaces in the patient, and environmental concerns. An inflow consisting of oxygen 500 ml/min and air 1000 ml/min – or somewhat different proportions – to a semiclosed circle absorber system combined with a narcotic and enflurane seems satisfactory for many clinical circumstances. The advent of agents that are metabolized less than those mentioned will make such an arrangement even more worthy of consideration.

John McIntyre, F.R.C.P.(C)
Department of Anaesthesia
University of Alberta Hospital
EDMONTON, Alberta