after peer review. Doctors Crone and Pavlin coauthored the article and many obstetric anaesthetists across the country made valuable contributions. We hope that we were able to reflect national peer opinion while avoiding delineation of rigid protocols based of necessity only on anecdotal case reports.

J.M. Davies MSc MD FRCPC
Department of Anaesthesia
Foothills Hospital
University of Calgary
Calgary, Alberta
S.K. Weeks MB BS FFARCS
Department of Anaesthesia
Royal Victoria Hospital
Montreal, Quebec

REPLY (2)

The comments by deMello and Restall (*Can J Anaesth* May 1990) about the use of laryngeal masks are timely. This device may indeed prove to be very useful in the management of failed intubation in obstetrics and elsewhere. However, experience with the laryngeal mask is not extensive and is limited mainly to the United Kingdom. The laryngeal mask is currently on trial in only one of the four North American teaching centres represented by the authors of our article. We await with interest the results of controlled studies.

Our article was not intended to provide a precisely detailed account of airway management techniques, including cricothyrotomy. We are grateful to Chung for his descriptions, but would suggest caution in recommending the use of an oesophageal obturator in every case, unless there are definite indications, e.g., gastric dilatation during trans-tracheal ventilation. In addition, the experience of obstetric anaesthetists would seem to contradict the opinion of Chung as to the practicality of IPPV with mask and cricoid pressure. Finally, the problem of fatigue during such a situation reinforces our recommendation for the presence of a trained assistant for the anaesthetist.

J.M. Davies MSc MD FRCPC
Department of Anaesthesia
Foothills Hospital
Calgary, Alberta
S.K. Weeks MB BS FFARCS
Department of Anaesthesia
Royal Victoria Hospital
Montreal, Quebec

REPLY (3)

We agree with Dr. Chung (*Can J Anaesth* 1990; 37: May) that all methods of obtaining access to the trachea may have problems in some patients. A preplanned algorithm for managing the airway is therefore necessary so as to proceed in a logical and timely way, through a series of techniques and approaches, until one proves successful. Of course in an emergency procedure where time is important, one may expect a significant incidence of complications including, unfortunately, death.

Specifically we attempt to pierce the cricothyroid membrane at an angle of approximately 30° such that the tip is caudad within the trachea. Where access is gained in a perpendicular

fashion, retraction of soft tissues of the neck cephalad has helped to place the needle in a better position. In patients with very large or swollen necks, use of a Tuohy needle has aided in maintaining patency. With experience, we have found that a #14 gauge Abbocath will suffice in most cases. While aspiration of oral contents is a potential problem, one study in dogs claimed no aspiration of oral water filled to 8 cm above the trachea if ventilatory rate via cricothyroid needle exceeded 40/minute. We have many times successfully ventilated a patient with this method while a definitive tracheostomy or indeed while oral endotracheal intubation was reattempted successfully. The efficacy of an oesophageal obturator has yet to be demonstrated.

Edward G. Pavlin MD
Associate Professor of Anesthesiology
University of Washington
Seattle, Washington, USA

Patient identity

To the Editor:

I happened to be looking up an article in the November 1989 edition of the Canadian Journal of Anaesthesia, when I came upon an article by Jan Davies *et al.* entitled "Difficult intubation in the parturient" (Vol 36: 668-674). I am writing to you because I feel that the identity of the patient in the article has not been adequately protected. An article on this subject from the New England Journal of Medicine¹ called "Unmasking the Lone Ranger" points out that putting sunglasses on Ronald Reagan or Michael Jackson does not disguise their identities. Similarly for the woman photographed in the article in the *Can J Anaesth* anyone who knows her could easily identify her because she has not been adequately disguised. I suppose that consent was received from this patient but I am sure she is not aware of the consequences of that consent and that her picture (in semi-clothed form) is now in the public record.

As researchers and educators, we are dependent upon the goodwill of the patients who kindly consent to participation in our studies and teaching exercises. In this article, I feel that protection of the confidentiality of this person has been breached.

Marsha M. Cohen MD FRCPC
Winnipeg

REFERENCE

- 1 Slue WE. Unmasking the lone ranger. *New Engl J Med* 1989; 321: 550-1.

REPLY

We concur with Cohen's premise that as physicians, educators, and researchers we are indebted to our patients who offer to