and oxygenation may be obtained with PTV and HFJV, we have some reservations about this technique in these patients. Access to the neck is often obstructed by various devices serving to stabilize the neck, rendering the placement of the transtracheal catheter problematic. These patients are often moved in the process of evaluation, as from stretcher to CT table, jeopardizing what may be already a tenuous airway. Finally, there is little evidence that the airway is adequately protected from soiling or aspiration in a patient population that is at significant risk for this event. Yealy cites two animal experimental models that suggest the risk of aspiration during PTV is low. However, in Yealy's own work, 66 per cent of dogs, at 45° of elevation, being ventilated with PTV and HFJV, suffered some degree of aspiration. There was no evidence of aspiration in animals intubated with a cuffed tracheal tube, under the same conditions. In the above situation, we are of the opinion that the priority is a secure and protected airway and that PTV with HFJV may be employed to ensure adequate ventilation and oxygenation while a plan is being formulated to secure the airway with a cuffed tube.

Airway management of patients with known or suspected cervical spinal injury or instability should be dictated by common sense and care and not by a dogmatic approach and reliance on any particular algorithm. This statement is supported by both the critical care literature and the accumulated clinical experience in the management of these patients, in many centres, with a variety of techniques.

E. Crosby MD FRCPC
A. Lui MD FRCPC
Ottawa Civic Hospital
University of Ottawa

Fatal embolism

To the Editor:

In his editorial, "Fatal Embolism,"¹ Dr. Matthews correctly indicates the need for "compulsory national reporting of (anaesthetic-related) mortality and serious morbidity." He suggests that this may be a role for the Department of Health and Welfare.

An alternative source of reporting of mortality, at least, are the provincial coroners and medical examiners. As they already review deaths, it would be logical to have them report to a central office. At the Annual Meeting of Chief Coroners and Medical Examiners (September, 1989, in Quebec City), we presented a systematic method for the investigation of anaesthetic-related deaths. Using such a common data base, information could be collected and collated. We suggested that, at a minimum, all intraoperative deaths become "notifiable," i.e., require mandatory reporting to the coroner/medical examiner. Better still would be a common period of 10–15 days postoperatively for notification of a death. However, the coroners and medical examiners operate under provin-

cial law, with twelve different acts. Currently, only three of the provinces/territories (Alberta, Manitoba, Prince Edward Island) require (by law) notification of operative deaths. Amendment of these acts to provide common times and conditions for reporting would be a major step towards helping us to know how many people die in Canada each year because of anaesthetic-related mishaps, the causes of these, and possible preventable factors.

J.M. Davies, MSc MD FRCPC
J.N. Armstrong MD
Department of Anaesthesia
Foothills Hospital at the University of Calgary

REFERENCE

- 1 Matthews RL. Fatal air embolism. *Can J Anaesth* 1990; 37: 12–4.

Prevention of obstruction of epidural catheter by blood clot

To the Editor:

I read with interest the case report described by Cohen and Amar.¹ In the described case it was possible to dislodge a blood clot by reinserting a new stylet wire through the epidural catheter. A simple measure is described which can prevent blood clot formation when an epidural catheter enters a blood vessel and is positive for blood on aspiration. A 10-ml syringe filled with saline is attached to the epidural catheter and is flushed while the catheter is partially withdrawn. This process is repeated until blood aspiration test is negative. This simple measure of flush and partial withdrawal helps the epidural catheter to come out of the vessel. We have not encountered accidental intravenous injection of local anaesthetic and very rarely need to resite the catheter in patients when an epidural vessel has been punctured.

Chandra M. Kumar FFARCS
Department of Anaesthesia
South Cleveland Hospital
Middlesbrough
Cleveland TS4 3BW

REFERENCES

- 1 Cohen S, Amar D. A simple solution to blood clot obstruction of epidural catheter during labour. *Can J Anaesth* 1990; 37: 143–4.