Neuromuscular blocking agents used with antibiotics

To the Editor:

Recently Dupuis *et al.* reported the action of atracurium and vecuronium in patients receiving aminoglycosides.¹ They showed no differences in onset time, but clinical duration and time to recovery were longer in patients receiving tobramycin or gentamicin and paralyzed with vecuronium than for controls, while the block produced by atracurium was not influenced by the presence of therapeutic serum levels of tobramycin or gentamicin. These findings are not surprising even though the anaesthesia literature has many case reports concerning aminoglycoside antibiotics contributing to the prolongation of non-depolarizing neuromuscular blocking agents. The authors stated in their paper that, "except for these isolated case reports, there are no clinical trials in man on the effects of aminoglycosides on neuromuscular block produced by muscle relaxants." A more thorough literature search, would have found the article, "Neuromuscular Blocking Effects of Tobramycin, Gentamicin and Cefazolin" by M. Lippman *et al.* in which 40 patients were studied.² We used d-tubocurarine as the muscle relaxant because the newer short-acting non-depolarizing relaxants were not available. Succinylcholine, used for intubation, was also monitored and studied (prior to the first dose of d-tubocurarine). Subsequent doses of d-tubocurarine were also studied and monitored.

The responses of the patients receiving tobramycin, gentamicin, or cefazolin did not differ from those of the control group. This was true when comparisons were made of the base-line neuromuscular data, duration of succinylcholine block, the potency, duration of block, recovery rate, train-of-four fade at 50 per cent block, tetanic trend, response to double stimuli, post-tetanic effect and reversibility of the block subsequently induced with d-tubocurarine. There were patients in each group who appeared to be resistant to d-tubocurarine. Not all patients receiving 0.2 mg · kg⁻¹ d-tubocurarine produced a block greater than 60 per cent. Two patients, one each in the tobramycin and cefazolin groups, required a third dose of 0.1 mg · kg⁻¹ of d-tubocurarine to produce 60 per cent suppression of the electromyograph. A third patient in the tobramycin group required a fourth dose. No problems were encountered in reversing the neuromuscular blockade following the administration of neostigmine, although the degree of neuromuscular transmission before neostigmine varied. In summary, gentamicin and tobramycin did not increase the sensitivity to d-tubocurarine nor did cefazolin.

Maurice Lippmann MD
Dora Hsu MD
Elaine Yang MD
Department of Anesthesiology
UCLA School of Medicine
Harbor/UCLA Medical Center Campus
1000 West Carson Street
Torrance, California 90509

REFERENCES

- 1 Dupuis JY, Martin R, Tetrault JP. Atracurium and vecuronium interaction with gentamicin and tobramycin. *Can J Anaesth* 1989; 36: 407-11.
- 2 Lippmann M, Yang E, Au E, Lee C. Neuromuscular blocking effects of tobramycin, gentamicin, and cefazolin. *Anesth Analg* 1982; 61: 767-70.

REPLY

We apologize for missing the above reference in our article.

Concerning the differences between the study of Lippmann *et al.*² and ours,¹ we would like to underline the following points of difference between the reports.

- 1 The neuromuscular relaxants studied were different. d-Tubocurarine was used in increasing dosages by Lippmann *et al.* while we used only intubating dosages of vecuronium and atracurium.
- 2 We did not use succinylcholine before d-tubocurarine.
- 3 The recovery index used by Lippmann *et al.* was the 25-50 per cent recovery time for twitch. We used the 0-25 per cent recovery time for twitch and the spontaneous recovery time for T4/T1 ratio to 0.70.
- 4 The EMG of the thumb with ulnar stimulation at 0.1 Hz, train-of-four stimuli at 2 Hz at "appropriate times," "paired stimuli 3 msec apart followed by a tetanic stimuli of 50 Hz of 5 seconds duration" were used by Lippmann *et al.* We used the EMG response of the hypothenar muscles and the ulnar nerve was stimulated at 0.05 Hz only in a train-of-four mode.
- 5 We did not use neostigmine to reverse the neuromuscular block.

All these differences could account for the different findings of the two studies and although we conclude that "vecuronium shows a significant increase in its duration in the presence of therapeutic serum levels of gentamicin or tobramycin while atracurium does not" it does not mean that vecuronium is not clinically safe in the presence of therapeutic serum levels of gentamicin or tobramycin.

René Martin MD
Department of Anaesthesia
University of Sherbrooke