FIGURE Changes in haematocrit (Hct), platelet count and prothrombin time (mean $\pm$ SEM). * $P < 0.05$ Student's paired t test. 0 = preoperative; P = postoperative.

unintentional, increase in almost every patient. The Hct decreased on the second postoperative day and returned to the baseline level six days later (Figure). Prothrombin time was prolonged and platelets were moderately reduced at the end of surgery and the following day (Figure). There were no relevant changes in the serum electrolytes at any time.

A few hours after the end of the surgical procedure, two patients experienced hypovolaemic shock, which was immediately corrected by infusion of fluids, bicarbonate, and calcium gluconate. Four infants were transfused in the postoperative period. There were no complications related to arterial catheterism nor perioperative deaths.

The majority of our patients (95 per cent) had to be transfused intraoperatively and the principal problems we experienced were connected with blood loss. In agreement with other authors,^{2,3} we recommend the use of short term arterial catheterization which allows blood-gas analysis, continuous measurement of blood pressure, and easy sampling for serial Hct determination. The last two variables together represent a reliable index for blood transfusion and fluid replacement in small children.⁴ In the postoperative period strict surveillance of vital signs and Hct is mandatory. Although coagulation disorders did not occur as a result of surgery in this study, monitoring of

coagulation indices is also necessary in large mass blood replacement.^{3,5}

R. Imberti MD
D. Locatelli* MD
M. Fanzio MD
N. Bonfanti* MD
I. Preseglio MD

Servizio di Anestesia e Rianimazione II –
IRCCS Policlinico S. Matteo and

*Dipartimento di Chirurgia,
Sezione di Neurochirurgia,
University of Pavia,
27100 Pavia – Italy.

REFERENCES

- 1 Kearney RA, Rosales IK, Howes WJ. Craniosynostosis: assessment of blood loss and transfusion practices. *Can J Anaesth* 1989; 36: 473–7.
- 2 Sholtes JL, Thauvoy Ch, Moulin D, Gribomont BF. Craniofaciosynostosis: anesthetic and perioperative management. *Acta Anaesthesiol Belg* 1985; 36: 174–85.
- 3 Vercauteren M, Van Vyve M, Janssen L, Hanegreets G, Selosse P. Craniostenosis: the importance of the anesthesiologist. *Acta Anaesthesiol Belg* 1985; 36: 168–73.
- 4 Steward DJ. *Manual of Pediatric Anesthesia*. 2nd ed, New York: Churchill Livingstone, 1985, 76.
- 5 Sette MP, Pietrini D, Bellani A, Bracali AM. Interventi per craniostenosi nelle prime epoche della vita. *Minerva Pediatr* 1983; 35: 159–5.

REPLY

We would like to thank Dr. Imberti and colleagues for providing more information on this infrequently addressed topic. Their figures for blood loss, though greater than in our study (28 per cent of estimated blood volume (EBV) overall) are in keeping with one other study.² The facts that 95 per cent of their patients required blood transfusions, nine per cent developed severe metabolic derangements secondary to blood loss, and four per cent developed hypovolaemic shock postoperatively, underline the importance of blood and fluid management of these patients.

Our study showed significant differences in blood loss between single suture craniectomies (sagittal or unicoronal) and multiple suture or metopic craniectomies. It would have been of interest to know this breakdown for their data. Sagittal craniectomies constitute the vast majority of cases, and blood loss during this procedure has generally been found to be less than 25 per cent of EBV^{1,3,4} (although in one study as high as 40 per cent of EBV²). The type of operative procedure may account for this difference as surgical approaches vary among institutions. We feel that patients with simple sagittal synostosis represent a lower risk group for perioperative complications and agree with others³ that invasive monitoring is not warranted in this group. For multiple suture craniectomies, invasive monitoring is advantageous.

For sagittal craniectomies, local practice should be evaluated to determine whether blood loss approximates the average or significantly exceeds it. Invasive monitoring may be warranted if blood loss is excessive.

Ramona A. Kearney MD FRCPC
Department of Anaesthesia
Dr. Charles A. Janeway Child Health Centre
St. John's, NF

REFERENCES

- 1 Kearney RA, Rosales JK, Howes WJ. Craniostomosis: assessment of blood loss and transfusion practices. *Can J Anaesth* 1989; 36: 473-7.
- 2 Scholtes JL, Tharvoy Ch, Moulin D, Gribomont BF. Craniofaciosynostosis: anaesthetic and perioperative management. *Acta Anaesthesiol Belg* 1985; 3: 176-85.
- 3 Diaz JH, Lockhart CH. Hypotensive anaesthesia for craniectomy in infancy. *Br J Anaesth* 1979; 51: 233-5.
- 4 Albright AL. Operative normalization of skull shape in sagittal synostosis. *Neurosurgery* 1985; 17: 329-31.

Triple lumen catheter: a potential hazard

To the Editor:

In many intensive care units (ICU), central triple lumen catheters are commonly used. Recently, in our institution, such a catheter (Baxter, model 703-K200H-7F) was inserted without incident into the right subclavian vein of a 45-year-old woman while she was in the ICU. Six days after its insertion, while the nurses were moving the patient from the supine to the semisitting position, one of them felt a little resistance. Almost immediately, they realized that the catheter had broken just proximal to the suture wing (Figure) and they immediately clamped the distal part of the catheter. When the house officer arrived, he cut the suture that anchored the catheter to the skin and removed the distal part. Because the problem was identified and treated quickly the patient did not experience any significant haemorrhage or air embolism.

We were concerned with two aspects of this incident: first, the nursing staff was certain that the catheter had been submitted only to moderate traction; second, the silk suture did not break before the catheter. Trying to recreate the same conditions, we sutured a new catheter to a table, stretched it with moderate traction and it broke at exactly the same spot as the first. Several other catheters from the same lot were tested and all presented the same problem. The suture wing of the Baxter 703-K200H-7F is glued by the manufacturer to the catheter.

The manufacturer was notified of the problem and the catheter was changed for one of another design. This

FIGURE Broken catheters.

incident did not have any serious consequences. However, with two 16 ga and one 18 ga lumen opened, serious haemorrhage or air embolism could easily have occurred.¹⁻²

Liette Isabel MD
Jean-Pierre Baribault MD FRCPC
Department of Anaesthesia
Hôpital Enfant-Jésus
Université Laval
1401, 18^e rue
Québec

REFERENCES

- 1 Cruz CA, Bedford RF. Complications of invasive cardiovascular monitoring. *Seminars in Anesthesia* 1990; 9: 16-25.
- 2 Kloosterboer TB, Springman SR, Coursin DB. Subclavian vein catheter as a source of air emboli in the sitting position. *Anesthesiology* 1986; 64: 411.

REPLY

We thank you for giving us the opportunity to comment.

From our review of the product complaint, we determined that the catheter separated at the tubing/suture sleeve junction. We were not advised of the amount of force applied either on the unit which separated or subsequently on those that were tested by the hospital. As part of quality assurance testing during manufacture these glue junctions are subjected to a controlled pull force - a force which is considered to exceed normal use forces. The four sets that were tested by the hospital were examined by our quality engineers and found to be within specifications.

As part of ongoing product development, and at the request of users, the fixed suture loop was changed to an optional suture loop on October 15, 1989. This loop change eliminated the glued junctions and gave the users a choice of where to place the loop. The new design that was referred to in the above letter