CORRECTIONS

SURGICAL STIMULATION DOES NOT ENHANCE VENTILATORY CHEMOREFLEXES DURING ENFLURANE ANAESTHESIA IN MAN

A.M. LAM, J.L. CLEMENT AND R.L. KNILL

Canadian Anaesthetists' Society Journal 27: 22-28 (January 1980)

Table I, page 25. In Group I, the Pa_{CO_2} for anaesthesia alone should be 7.8 ± 0.4 kPa and the value for anaesthesia and surgery should be 7.4 ± 1.1 kPa. These values differ significantly from each other ($p < .05$).

REDUCTION OF POST-OPERATIVE VOMITING BY PREOPERATIVE ADMINISTRATION OF ORAL METOCLOPRAMIDE

M.J. DIAMOND AND M. KEERI-SZANTO

Can. Anaes. Soc. Jour. 27: 36 (January 1980)

TABLE VI
51 ABDOMINAL HYSTERECTOMIES

	Vomited	Did not vomit
Control	13	13
Treatment	8	17

The reduction in postoperative vomiting in the treatment group does not reach statistical significance.

NON-KETOTIC HYPERGLYCAEMIAS

K.G. BELANI AND K.A. SAMI

Can. Anaes. Soc. Jour. 27: 39 (January 1980)

The name of K.G. Belani, M.D., should be added to the signatures on this letter.