FIGURE 1 The faucial pillars, soft palate, and uvula are easily seen.

FIGURE 2 The distance from thyroid notch to lower border of the mandible is less than 6.0 cm.

of Patil *et al.*² Those authors found this measurement to predict patients at risk for difficult visualization.

Successful direct visualization of the larynx and tracheal intubation requires alignment of the axes of mouth, pharynx, larynx³ and deformation of the soft tissues in the floor of the mouth. The method of Mallampati would seem to predict the patient at risk for difficult intubation by assessing the amount of soft tissue that would be deformed during laryngoscopy. The method of Patil would seem to predict the patient at risk for difficult laryngoscopy by indirectly assessing the size of the region to which the soft tissue has to be deformed. Perhaps a short mandibular to thyroid distance should be added to those list of anatomic signals that herald a difficult intubation. Inability to visualize the uvula and faucial pillars should be taken as strong evidence of a potentially difficult intubation; however, visualization of those same structures may not predict ease in exposing the larynx by direct laryngoscopy.

Curtis L. Baysinger MD
William Evans DO
Robert Middaugh MD
John Culclasure MD
Douglas Culling DO
Anesthesia and Operative Service
Brooke Army Medical Center
Fort Sam Houston, Texas 78234

REFERENCES

- 1 Mallampati SR, Gatt SP, Gugino LD *et al.* A clinical sign to predict difficult tracheal intubation: a prospective study. *Can Anaesth Soc J* 1985; 32: 429-33.
- 2 Patil VU, Stehling LC, Zauder HL. Predicting the difficulty of intubation using an intubation gauge. *Anesthesiology Review* 1983; 10: 32-3.
- 3 Cass NM, James NR, Lines V. Difficult direct laryngoscopy complicating intubation for anaesthesia. *Br Med J* 1956; 1: 488-9.

REPLY

I would like to thank Baysinger *et al.* for their thought-provoking letter indicating that they had failed to expose the glottis in two patients in whom the faucial pillars, the soft palate and uvula could easily be visualized. These patients apparently had no airway pathology and the mobility of temporomandibular joint and cervical spine was said to be unrestricted. Although it is somewhat difficult to evaluate the situation with objectivity and pinpoint the cause of their difficulty, I would suspect that the use of tubocurarine prior to the administration of succinylcholine was at least, in part, responsible for the difficulty. With the growing popularity of using a nondepolarizer such as tubocurarine prior to succinylcholine, the efficacy of the latter is curtailed,¹⁻³ often leading to inadequate muscle relaxation. As a consequence, it becomes difficult to bring about the best possible "sniffing position" (flexion of the cervical spine and extension at the atlanto-occipital joint) and axes alignment.

In my experience as well as in the experience of others,⁴ failure to bring about optimal axes alignment is perhaps the most common cause of difficult orotracheal intubation. This type of situation frequently leads us to administer multiple doses of muscle relaxants and repeated attempts at intubation which, by virtue of haemorrhage and oedema, can only compound the difficulties. It is for this reason that no precurarization was employed in our study.⁵ Furthermore, the entire group of patients were subjected to the same pharmacologic doses and physical conditions. With such uniformity, there were no "inadequate exposures" of glottis in patients in whom faucial pillars, soft palate and uvula were visible. Nevertheless, it is still possible that in an occasional patient there may be anatomical variations in the mandible or cervical spine that are clinically subtle but would render axes alignment difficult. As our experience grows, we will be able to refine the existing clinical method and perhaps design some new methods to complement the former such that we can identify nearly all patients who are at risk for difficult orotracheal intubation.

As for Patil's method⁶ of predicting difficult intubation, it would seem that the larynx is much closer to the base of the tongue with a shorter distance between the lower border of the mandible and thyroid notch. It is not clear how often and in what

degree this difficulty occurs with a distance of ≤ 6.0 cm. In order to evaluate objectively, it is necessary to conduct a large-scale study.

S. Rao Mallampati MD
Department of Anesthesia
Harvard Medical School and
Brigham and Women's Hospital
75 Francis Street
Boston, Massachusetts
02115

REFERENCES

- 1 Cullen DJ. The effect of pretreatment with nondepolarizing muscle relaxants on the neuromuscular blocking action of succinylcholine. *Anesthesiology* 1971; 35: 572-8.
- 2 Freund FG, Rubin AP. The need for additional succinylcholine after d-tubocurarine. *Anesthesiology* 1972; 36: 185-7.
- 3 Walls LF, Dillon JB. Clinical studies of the interaction between d-tubocurarine and succinylcholine. *Anesthesiology* 1969; 31: 39-44.
- 4 Salem MR, Mathrubutham M, Bennett EJ. Difficult intubation. *N Engl J Med* 1976; 295: 879-81.
- 5 Mallampati SR, Gatt SP, Gugino LD *et al.* A clinical sign to predict difficult tracheal intubation: a prospective study. *Can Anaesth Soc J* 1985; 32: 429-34.
- 6 Patil VU, Stehling LC, Zauder HL. Predicting the difficulty of intubation utilizing an intubation gauge. *Anesthesiology Review* 1983; 10: 32-3.

Combined subarachnoid and epidural block for Caesarean section

To the Editor:

I would be grateful for the use of your columns to comment on the letter from Dr. Rawal.¹ The single-space double-needle technique has been described previously^{2,3} and I would like to mention our modification of the technique.⁴ This involves clamping the spinal needle with a light-weight artery clip at the point where it enters the Tuohy needle. This avoids the difficulty commonly encountered with a long flexible needle, whereby when attempting to firmly attach a syringe with a leak-proof seal and then inject, the needle may be advanced through the ventral surface of the dura, or passed out through the original dorsal puncture. In either case, the local anaesthetic solution is deposited extradurally, resulting in failure of the technique.

The dose and drug recommended by Rawal was 1.5 to 2.0 ml of hyperbaric 0.5 per cent bupivacaine. In their 24 cases they noted that the block rose to T₄ with the subarachnoid injection in only 25 per cent of the patients.

We use 1.5 to 1.6 ml of 0.5 per cent isobaric (slightly hypobaric) bupivacaine and, in a series of over 400

patients, the subarachnoid injection consistently reached T₂ to T₄. In less than five per cent of patients does the epidural cannula need to be used to bring the sensory block up to a level suitable for surgery. This has the advantage of reliably quick onset, because the patient can be moved into theatre immediately after the block is performed and preparation for operation commenced. We now, except in the most life-threatening emergencies, use this technique for emergency Caesarean section, as the patient, with a skilled team, can be in the operating room with a block at T₅ and undergoing skin preparation, five minutes after starting the procedure.

At the beginning of our series, we noted a similar incidence of hypotension to that noted by Rawal, i.e., about 10 to 15 per cent. We now administer 9 mg ephedrine IV immediately after the epidural cannula is inserted, plus supplementary injections of 6 mg if the systolic blood pressure falls below 110, following a "pre-load" of two litres of lactated Ringer's solution. The incidence of hypotension, defined as systolic blood pressure below 100, is now less than five per cent.

I would agree with Dr. Rawal about the incidence of headache. We had two spinal headaches in the first 400 patients. Whether or not this is attributable to extradural opiates is a matter of controversy. We use 0.3 mg buprenorphine in 10 ml 0.25 per cent bupivacaine extradurally and the patients are allowed to mobilize as soon as they have sufficient motor power.

A final point to make is that although our sensory block is usually at a high thoracic level, we have had no respiratory problems from intercostal weakness, suggesting again the beneficial effect of 0.5 per cent isobaric bupivacaine in producing a differential marked sensory block with little motor block.

Brian Dennison
Department of Anaesthesia
South Cleveland Hospital
Marton Road
Middlesbrough
Cleveland
TS3 4BW United Kingdom

REFERENCES

- 1 Rawal N. Single segment combined subarachnoid and epidural block for Caesarean section. *Can Anaesth Soc J* 1986; 33: 254-5.
- 2 Coates MB. Combined sub-arachnoid and epidural techniques. *Anaesthesia* 1982; 37: 87-9.
- 3 Muntaz MH, Daz M, Juz M. Another single space technique for orthopaedic surgery. *Anaesthesia* 1982; 37: 90.
- 4 Nickalls RW, Dennison B. A modification of the combined spinal and epidural technique. *Anaesthesia* 1984; 39: 935-6.