REFERENCES

- 1 Virtue R, Sherrill DL, Swanson GD. Uptake of nitrous oxide by man. *Can Anaesth Soc J* 1982; 29: 424-7.
- 2 Severinghaus JW. Rate of uptake of nitrous oxide in man. *J Clin Invest* 1954; 33: 1183-9.
- 3 Aldrete JA, Lowe HJ, Virtue RW. Low flow and closed system anesthesia. New York, Grune and Stratton, 1979.
- 4 Lowe HJ, Ernst EA. The quantitative practice of anesthesia. Baltimore, William and Wilkins, chapter 5 pp 67-97, 1981.
- 5 Lin CY. Nitrous oxide uptake in adults (Abstract). *Anesthesiology* 1982; 57: S372.
- 6 Epstein RM, Rackow H, Salanitro E, et al. Influence of the concentration effect on the uptake anesthetic mixture: The second gas effect. *Anesthesiology* 1964; 25: 364-71.
- 7 Stoelting RK, Eger EI, II. An additional explanation for the second gas effect. A concentrating effect. *Anesthesiology* 1969; 30: 273-7.
- 8 Fink BR. Diffusion anoxia. *Anesthesiology* 1955; 16: 511-9.
- 9 Lin CY, Mostert JW. Inspired oxygen and nitrous oxide concentration in essentially closed circuits. *Anaesthetist* 1977; 26: 514-7.
- 10 Lin CY, Mostert JW, Benson DW. Closed circuit systems. A new direction in the practice of anesthesia. *Acta Anaesthesiol Scand* 1980; 24: 354-61.

REPLY

Dr. Lin and co-workers, ourselves and others, have for several years been trying to emphasize the tremendous waste and lack of scientific acumen in the general use of greater quantities of anaesthetic agents than the amount required by the patient. Recently, Dr. Lin has been especially vigorous in motivating the use of closed circle systems.^{1,2} We are entirely supportive of the practical concepts that he suggests for administration of nitrous oxide in the clinical setting.

However, scientifically we differ from the Chicago group. The purpose of our paper³ was to re-examine critically nitrous oxide uptake in man, using modern instrumentation and methods. Our instrumentation system consists of a specially designed miniature quadrupole mass spectrometer and a specially designed hot-wire anemometer flowmeter.⁴ The ionization chamber of the mass spectrometer is connected directly to a 35 ml flow-through tube, eliminating the traditional mass spectrometer sampling catheter. The response time of this system is less than 30 ms with a delay time of less than 1 ms for nitrous oxide. The inlet to the ionization chamber

and the hot-wire flowmeter are at the same physical location in the middle of the tube. Thus, there is no phase delay between the flow signal and the gas fraction signal. An on-line computer processes these signals directly every 20 ms to yield breath-to-breath gas exchange data.

Our method⁵ for estimating alveolar gas exchange (gas exchange from the lung to blood) utilizes a version of the Auchincloss algorithm,⁶ which is also used by other groups.⁷ It determines the uptake of the mouth and subtracts an estimate of nitrous oxide taken into the pulmonary stores (FRC) using approximately one per cent argon as a tracer gas. We and others have carried out an extensive analysis of these methods.^{7,8}

Dr. Lin has suggested that our "nitrous oxide experiment was carried out separately from the FRC measurement." However, this was not the case. The subjects, breathing room air, were switched onto the oxygen nitrous oxide mixture at time zero. Nitrogen wash-out was then measured simultaneously with the nitrous oxide wash-in, while the room air argon concentration was maintained. Real time measurements were combined off-line to yield our nitrous oxide uptake estimates.

These methods and our instrumentation system allow us to make the quantitative measurements shown in our paper. Furthermore, we have also quantitatively demonstrated the concentration and second gas effect, using nitrous oxide as the soluble gas.⁹

In contrast, the Chicago group uses simpler methods and qualitative arguments to motivate concepts that may be useful clinically, but are questionable from a scientific point of view. Consider their concepts from this perspective:

1. The Chicago group emphasizes the "mean constancy" of nitrous oxide uptake by considering strip chart recordings of nitrous oxide concentrations on linear scales. In contrast, we have calculated the nitrous oxide uptake and plot the data on log-log scales to emphasize small changes and the similarity to the predictive square root of time empirical model.

2. The Chicago group indicates they do not observe the concentration and second gas effects in their clinical measurement setting. However, they have not analyzed their strip chart recordings quantitatively for such an effect. Furthermore, they argue on a conceptual basis that these effects have been overemphasized because investigators have not considered FRC wash-in. However, the observation of these effects cannot be dismissed on this basis, since they can be demonstrated easily, using nitrous oxide as the soluble gas.⁹

3. The Chicago group has suggested that an alveolar-capillary diffusion barrier exists^{1,2} to account for their observations and concepts concerning nitrous oxide uptake. However, nitrous oxide has been used for a tracer gas in a variety of studies to estimate pulmonary blood flow.^{10,11} Diffusion limitations to uptake are not evident.

Finally, Dr. Lin objects to our terminology. Our use of "total uptake" for the uptake at the mouth emphasizes the uptake that the practicing anaesthesiologist must be concerned about — that which flows from the anaesthetic

machine. Furthermore, our use of "alveolar uptake" is consistent with the physiology literature.

George D. Swanson PhD
Robert W. Virtue MD PhD
Duane Sherrill MS
Department of Anesthesiology
University of Colorado Health Sciences Centre
Denver, Colorado, 80262

REFERENCES

- 1 Lin CY, Mostert JW. Inspired oxygen and nitrous oxide concentrations in essentially closed circuits. *Anesthetist*. 1977; 26: 514-7.
- 2 Lun CY, Mostert JW, Benson DW. Closed circuit system: a new direction in the practice of anaesthesia. *Acta Anaesthesiol Scand* 1980; 24: 354-61.
- 3 Virtue R, Sherrill DL, Swanson GD. Uptake of nitrous oxide by man. *Can Anaesth Soc J* 1982; 29: 424-7.
- 4 Sodal IE, Swanson GD, Micco AJ, Sprague E, Ellis DG. A computerized mass spectrometer and flow meter system for respiratory gas measurements. *Ann Biomed Eng* 1982; 10.
- 5 Swanson GD. Breath-to-breath considerations for gas exchange kinetics in *Exercise and bioenergetics and Gas Exchange*, Eds. Cerretelli P, Whipp BJ. p. 211-22. Elsevier/North Holland, Amsterdam, 1980.
- 6 Auchincloss JH, Gilbert R, Baude GH. Effect of ventilation of oxygen transfer during early exercise. *J Appl Physiol* 1966; 21: 810-8.
- 7 Beaver WL, Lamarra N, Wasserman K. Breath-by-breath measurement of true alveolar gas exchange. *J Appl Physiol* 1981; 51: 1662-75.
- 8 Swanson GD, Sodal IE, Reeves JT. Sensitivity of breath-to-breath gas exchange measurements to expiratory flow errors. *IEEE Trans Biomed Eng* 1981; 28: 749-54.
- 9 Johnson TS, Swanson GD, Sodal IE, Reeves JT, Virtue RW. A closed lung system study of inert gas absorption. *J Appl Physiol* 1979; 47: 240-4.
- 10 Cander L, Forester RE. Determination of pulmonary perenchymal tissue volume and pulmonary capillary blood flow in man. *J Appl Physiol* 1959; 14: 541-51.
- 11 Brady JP, Rigatto H. Pulmonary capillary flow in infants. *Circulation (Suppl. III)* 1969; 40: 50.

Renal haemodynamics with fentanyl

To the Editor

We read the paper "Effects of high dose fentanyl on renal haemodynamics in conscious dogs"¹ with interest as we published similar work on the effect of fentanyl 25 $\mu\text{g}\cdot\text{kg}^{-1}$ administered to the lightly anaesthetised normocapnic dog in the *British Journal of Anaesthesia*.² In our cardiovascularly stable animal model we injected fentanyl 25 $\mu\text{g}\cdot\text{kg}^{-1}$ over a ten-minute period into a peripheral vein and also found a statistically significant fall in mean arterial pressure (MAP) lasting for up to 90 min, i.e., longer than the effect noted by Dr. Priano in conscious dogs given the same dose. In contrast to Dr. Priano, however, the anaesthetised animals in our series also had a significant fall in estimated renal plasma flow (ERPF) after administration of fentanyl 25 $\mu\text{g}\cdot\text{kg}^{-1}$ (measured by PAH clearance), which was more marked than the changes noted by Bidwai *et al.*³ The fall in renal ERPF was also accompanied by a significant fall in urine volume (V) in our series; a parameter not measured in the conscious dog. In addition, we found a marked rise in renal vascular resistance (RVR) as the fall in ERPF was greater than the fall in MAP. Because of the similarity between our work and his own we feel that it would have been appropriate for Dr. Priano to have compared and contrasted the two in his paper. This example highlights a situation well recognised in Great Britain amongst anaesthetists: why do so many North American publications fail to acknowledge published British work on similar subjects?

Dr. Jennifer M. Hunter
Dr. R.S. Jones
The University Department of Anaesthesia
Royal Liverpool Hospital
Liverpool, L69 2BX, England

REFERENCES

- 1 Priano LL. Effects of high-dose fentanyl on renal haemodynamics in conscious dogs. *Can Anaesth Soc J* 1983; 30: 10-18.