REFERENCES

- 1 Kirsch JR, Diring MN, Borel CO, et al. Preoperative lumbar epidural morphine improves postoperative analgesia and ventilatory function after transsternal thymectomy in patients with myasthenia gravis. *Crit Care Med* 1991; 19: 1474-9.
- 2 Shulman M, Sandler AN, Bradley JW, et al. Post thoracotomy pain and pulmonary function following epidural and systemic morphine. *Anesthesiology* 1984; 61: 569-75.
- 3 Yeager M, Glass D, Neff R, et al. Epidural anesthesia and analgesia in high-risk surgical patients. *Anesthesiology* 1987; 66: 729-36.
- 4 Shenaq S. Con: epidural anesthesia is not a valuable adjunct to general anesthesia for abdominal vascular surgery. *Cardiothorac Anesth* 1989; 3: 509-12.

Perioperative stroke

To the Editor:

While reading DHW Wong's extensive and informative two-part review "Perioperative Stroke,"^{1,2} I came across a statement in Part I which troubled me, not because of what was said but by what was implied: "Halothane produces cerebral vasodilation with an increase in cerebral blood flow and increased intracranial pressure. On the other hand, isoflurane reduces cerebral oxygen consumption, preserves the normal energy state, and has minimal effect on intracranial pressure."²²⁸⁻²³³

In fact, things are not so clear as this. Dr. Michenfelder (Dr. Wong's reference 231), concluded that reasons for the apparent benefit from isoflurane in carotid endarterectomy were not as obvious as this. While most authors agree that halothane increases cerebral blood flow more than isoflurane, agreement on their effects on intracranial pressure is not as universal. For example, Kaieda *et al.*,³ in a rabbit model with a cryogenic cortical injury, were unable to demonstrate any difference in intracranial pressure at 1 MAC of isoflurane or halothane, or an equal dose of pentobarbital. Mutch *et al.*⁴ in a similar model, found no difference in intracranial pressure until mean arterial pressure had been increased 60% above baseline with phenylephrine.

Particularly given the difference in distribution of cerebral blood flow effected by the two agents,⁵ I believe that Todd and Drummond are correct when they conclude that "the simple fact that isoflurane produced consistent dose-related increases in intracranial pressure ... indicates that it should be used with caution in situations where intracranial compliance is compromised..."⁶

Ian J. Gilmour MD
University of Minnesota

REFERENCES

- 1 Wong DHW. Perioperative stroke. Part I: General surgery, carotid artery disease, and carotid endarterectomy. *Can J Anaesth* 1991; 38: 347-73.
- 2 Wong DHW. Perioperative stroke. Part II: Cardiac surgery and cardiogenic embolic stroke. *Can J Anaesth* 1991; 38: 471-88.
- 3 Kaieda R, Todd MM, Weeks JB, Warner DS. A comparison of the effects of halothane, isoflurane and pentobarbital anesthesia on intracranial pressure and cerebral edema formation following brain injury in rabbits. *Anesthesiology* 1989; 71: 571-9.
- 4 Mutch WAC, Patel PM, Ruta TS. A comparison of the cerebral pressure-flow relationship for halothane and isoflurane at haemodynamically equivalent end-tidal concentrations in the rabbit. *Can J Anaesth* 1990; 37: 223-30.
- 5 Hansen TD, Warner DS, Todd MM, et al. Distribution of cerebral blood flow during halothane versus isoflurane anesthesia in rats. *Anesthesiology* 1988; 69: 332-7.
- 6 Todd MM, Drummond JC. A comparison of the cerebral, vascular and metabolic effects of halothane and isoflurane in the cat. *Anesthesiology* 1984; 60: 276-82.

REPLY

Dr. Gilmour was quite correct in pointing out that the observed effects of halothane and isoflurane on intracranial pressure (ICP) are not uniform. In neurosurgical patients, for example, isoflurane anaesthesia has been shown to produce no significant effect on ICP.¹ In animal studies, when compared with equipotent concentrations of halothane, isoflurane produces lower^{2,3} or similar ICP levels.^{4,7} The observed differences may be due to differences in species studied, experimental design, instrumentation, supplementary anaesthetic drugs, and concomitant drug therapy. Furthermore differences in MAP and PaCO₂, presence or absence and type of ischaemic injury may have overriding effects. ICP changes are important as they directly influence cerebral perfusion pressure (CPP), which equals the difference between mean arterial pressure (MAP) and ICP.

Todd and Drummond⁵ showed that in cats, halothane produced greater increases in CBF and less decreases in CMRO₂ than isoflurane. In spite of the differences in CBF, the ICP changes produced by the two agents were similar. Since ICP changes should follow the changes in cerebral blood volume (CBV), there appears to be a dissociation between the effects of anaesthetics on CBF and CBV. One explanation is that the study animals with no intracranial pathology may have high cerebral compliance.^{6,8} Another explanation is that halothane affects the inflow and outflow resistance equally, whereas isoflurane has less effect on the arterioles and even less effect on the outflow resistance, thus allowing accommodation of a larger CBV and resulting in higher ICP at lower CBF.^{4,5} It is also possible, since the xenon clearance method used measures, predominantly, cortical CBF, that the observed differences in CBF were not representative of whole-brain differences.⁹

It has been suggested that the control of CPP may not be crucial during light isoflurane anaesthesia, as the uncoupling of local flow and metabolism by isoflurane is dose-dependent.¹⁰⁻¹² Isoflurane produces less cerebral vasodilation compared to enflurane and halothane, and might therefore cause a relatively

less unfavourable redistribution of flow during incomplete regional ischaemia.¹³ For situations where intracranial compliance is compromised, deep isoflurane should be avoided. Thiopentone would be a more appropriate agent where brain protection is required.

David H.W. Wong MB BS FRCPC
Department of Anaesthesia
University of British Columbia
Vancouver

REFERENCES

- 1 Campkin TV. Isoflurane and cranial extradural pressure. A study in neurosurgical patients. *Br J Anaesth* 1984; 56: 1083-7.
- 2 Artru AA. Relationship between cerebral blood flow and CSF pressure during anesthesia with halothane or enflurane in dogs. *Anesthesiology* 1983; 58: 533-9.
- 3 Artru AA. Relationship between cerebral blood volume and CSF pressure during anesthesia with isoflurane or fentanyl in dogs. *Anesthesiology* 1984; 60: 575-9.
- 4 Mutch WAC, Patel PM, Ruta TS. A comparison of the cerebral pressure-flow relationship for halothane and isoflurane at haemodynamically equivalent end-tidal concentrations in the rabbit. *Can J Anaesth* 1990; 37: 223-30.
- 5 Todd MM, Drummond JC. A comparison of the cerebrovascular and metabolic effects of halothane and isoflurane in the cat. *Anesthesiology* 1984; 60: 276-82.
- 6 Drummond JC, Todd MM. The response of the feline cerebral circulation to PaCO₂ during anesthesia with isoflurane and halothane and during sedation with nitrous oxide. *Anesthesiology* 1985; 62: 268-73.
- 7 Kaieda R, Todd MM, Weeks JB, Warner DS. A comparison of the effects of halothane, isoflurane, and pentobarbital anesthesia on intracranial pressure and cerebral edema formation following brain injury in rabbits. *Anesthesiology* 1989; 71: 571-9.
- 8 Brüssel T, Fitch W, Brodner G, Arendt I, Van Aken H. Effects of halothane in low concentrations on cerebral blood flow, cerebral metabolism, and cerebrovascular autoregulation in the baboon. *Anesth Analg* 1991; 73: 758-64.
- 9 Hansen TD, Warner DS, Todd MM, Vus LJ, Trawick DC. Distribution of cerebral blood flow during halothane versus isoflurane anesthesia in rats. *Anesthesiology* 1988; 69: 332-7.
- 10 Hoffman WE, Edelman G, Kochs E, Werner C, Segil L, Albrecht RF. Cerebral autoregulation in awake versus isoflurane-anesthetized rats. *Anesth Analg* 1991; 73: 753-7.
- 11 Maekawa T, Tommasino C, Shapiro HM, Keifer-Goodman J, Kohlenberger RW. Local cerebral blood flow and glucose utilization during isoflurane anesthesia in the rat. *Anesthesiology* 1986; 65: 144-51.
- 12 McPherson RW, Traystman RJ. Effects of isoflurane on cerebral autoregulation in dogs. *Anesthesiology* 1988; 69: 493-9.
- 13 Michenfelder JD, Sundt TM, Fode N, Sharbrough FW. Isoflurane when compared to enflurane and halothane decreases the frequency of cerebral ischemia during carotid endarterectomy. *Anesthesiology* 1987; 67: 336-40.