cation consisting of intramuscular morphine 5 mg and promethazine 25 mg was given at 0855 hours. General anaesthesia was induced at 1020 hours, maintained with isoflurane, plus fentanyl which was given in increments to a total of 300 µg. Following the operation, the orthopaedic surgeon gave at 1230 hours 5 ml (2.5 mg) of a preservative-free morphine solution through a small catheter 4–5 centimeters above the laminotomy site outside the dura. The various layers were closed. This direct application of morphine over the dura to effect postoperative analgesia following spinal operations, has recently been described¹ and forms part of a standard protocol by our orthopaedic surgeons.

No analgesics were given in the recovery room. Following discharge to the ward at 1415 hours the patient was monitored for signs of respiratory depression. At 1730 he was found to be somnolent, cyanotic and breathing seven times/minute. Arterial blood gas analysis (FI_{O_2} 0.21) revealed: PO_2 29 mmHg, PCO_2 of 56 mmHg. Oxygen was then administered. The respiratory rate fell to four/minute. Naloxone 0.2 mg was given intravenously over several minutes. The patient became alert and the respiratory rate increased to ten/minute. Repeat arterial blood gas analysis revealed: PO_2 122 mmHg, PCO_2 51 mmHg. Two hours later, the respiratory rate again decreased to six breaths/minute. Naloxone 0.1 mg was injected intravenously at hourly intervals until the following morning. The postoperative course thereafter was uneventful.

We report this interesting case to highlight the vast range of susceptibility to the respiratory depressant effects of epidurally administered narcotics, in this case given by the surgeon. Although not apparent at the time of operation, an inadvertently created opening in the dura cannot be definitely ruled out, allowing direct entry into the subarachnoid space of a portion of the morphine-containing solution. The morphine concentration in the CSF might therefore have been significantly higher than usual. This case supports the need for close postoperative observation of such patients in order to avoid a tragedy.

Christopher Jyu MD
Jonathan D. Lamb MD FRCP(C)
Department of Anaesthesia
University Hospital
Saskatoon, Saskatchewan

REFERENCE

- 1 *Rechtine GR, Reinert CM, Bohlman NH.* The use of epidural morphine to decrease post operative pain in patients undergoing lumbar laminectomy. *J Bone and Joint Surg* 1984; 66: 113–6.

Erratum

Re: *Palazzo, M.G.A., Strunin, L.* Anaesthesia and emesis. II. Prevention and management. *Can Anaesth Soc J* 1984; 31: 407–416 (July).

The sentence (page 413, line 27, right hand column) "Our drug of choice for prophylaxis would be droperidol 0.175 mg·kg⁻¹ intravenously, given preoperatively, although a smaller dose can be effective."

should read

"Our drug of choice for prophylaxis would be droperidol 0.0175 mg·kg⁻¹ intravenously, given preoperatively, although a smaller dose can be effective."