Correspondence

Serum potassium change during the TURP syndrome by cell volume regulation

To the Editor:

In the TURP syndrome, changes in serum potassium have not been evaluated precisely. Jensen¹ described the causes of potassium change during the TURP syndrome, but others should be added.

If the serum potassium concentration does not change during the TURP syndrome it suggests that intracellular potassium is extruded to the extracellular space, to compensate for renal potassium losses and keep serum potassium constant.

Non-electrolyte hypotonic solutions, glycine, sorbitol or mannitol, are usually used as a perfusate for TURP, and the large leakage into the blood vessels makes the extracellular fluid hypotonic. Hypotonic stress in extracellular space induces cell swelling, but the swollen cells tend to return to their original volume within a few minutes.^{2,3} This physiological cell volume "regulatory volume decrease (RVD)," involves efflux of potassium and water from the intracellular space.^{4,5}

Potassium extrusions from the cells by RVD have been reported in erythrocytes, lymphocytes, hepatocytes, and renal cells.²⁻⁴ Physiologically, these cells may be exposed to hypotonic stress by water drinking but they should keep their cell volume constant. In the TURP syndrome, these cells may also show an RVD response and extrude potassium into the extracellular space.

The RVD response should be considered in evaluating changes in serum potassium concentration during TURP syndrome.

Munetaka Hirose MD
Yoshifumi Tanaka MD
Department of Anaesthesia
Kyoto Prefectural University of Medicine
Kamigyoku Kyoto 602
Japan

REFERENCES

- 1 Jensen V. The TURP syndrome. *Can J Anaesth* 1991; 38: 90-7.
- 2 Haddad P, Graf J. Volume-regulatory K⁺ fluxes in the isolated perfused rat liver: characterization by ion transport inhibitors. *Am J Physiol* 1989; 257: G357-63.
- 3 Joannidis M, Volkl H, Pfaller W, Lang F. Volume-regulatory potassium release from isolated perfused rat kidney. *Renal Physiol Biochem* 1989; 12: 338-46.
- 4 Eveloff JL, Warnock DG. Activation of ion transport systems during cell volume regulation. *Am J Physiol* 1987; 252 F1-10.
- 5 Chamberlin ME, Strange K. Anisotonic cell volume regulation: a comparative view. *Am J Physiol* 1989; 257: C159-73.

REPLY

I appreciated the discussion concerning serum potassium change during TURP which was contributed by Dr. Hirose and Dr. Tanaka. Potassium is mostly an intracellular ion. When serum [K⁺] is diluted by irrigating fluid, then intracellular potassium is extruded to the extracellular space in compensation. As well explained by Dr. Hirose and Dr. Tanaka, the regulatory volume decrease (RVD) response would help to explain this potassium flux, so there would be no serum [K⁺] change during relatively small absorptions. The distribution of potassium in the body is regulated by a myriad of factors. The normal high intracellular to extracellular potassium concentration is dependent on the Na-K-ATPase pump. Other influences on the distribution of potassium between the intracellular and extracellular fluid compartments are three hormones (insulin, epinephrine, and aldosterone), acid-base status, and plasma tonicity.¹ The effect of tonicity is opposite to that of the RVD response.

Serum potassium change during TURP with glycine, 1.5%, depends on the rate and amount of irrigating fluid absorption. If 300 ml of irrigating fluid is absorbed during the case, then there is no change in serum [K⁺].² If a patient absorbs >300 ml of irrigating fluid intravascularly, then serum [K⁺] is increased by 0.5 mmol · L⁻¹ by the end of the case.³ An increased serum [K⁺] at the end of the surgery is a sign of irrigating fluid absorption;³ subsequently, a decrease in serum [K⁺] occurs 6 to 48 hr after the surgery.⁴

After glycine is absorbed, it moves intracellularly where it is metabolized. The system for glycine transport across the cell membrane is connected to the Na-K-ATPase pump.⁴ Norlen states that a high plasma concentration of glycine may interfere with the transmembrane electrolyte exchange, leading to a rise in serum [K⁺].⁴ The decrease in serum [K⁺] which occurs six hours after surgery may indicate a shift of potassium intracellularly, after the glycine has been metabolized. Transient acidosis, leakage from injured prostatic cells, the RVD response and haemolysis might also increase the serum [K⁺] at the end of the TURP.

V. Jensen MD FRCPC
Edmonton

REFERENCES

- 1 Smith JD, Bia MJ, DeFronzo RA. Clinical disorders of potassium metabolism. In: Arieff AI, DeFronzo RA (Eds.). *Fluid, Electrolyte and Acid-Base Disorders*. New York: Churchill Livingstone Inc. 1985: 413-509.
- 2 Sellevold O, Breivik H, Tveter K. Changes in oncotic pressure, osmolality and electrolytes following transurethral resection of the prostate using glycine as irrigating solution. *Scand J Urol Nephrol* 1983; 17: 31-6.
- 3 Hahn RG, Berlin T, Lewenhaupt A. Factors influencing the osmolality and the concentrations of blood haemoglobin and electrolytes during transurethral resection of the prostate. *Acta Anaesthesiol Scand* 1987; 31: 601-7.
- 4 Norten H, Allgen LG, Vinnars E, Bedrelidou-Classon G. Glycine solution as an irrigating agent during transurethral prostatic resection. *Scand J Urol Nephrol* 1986; 20: 19-26.

Cuff tears: can they be prevented?

To the Editor:

The nasotracheal route is usually utilized in patients selected for awake tracheal intubation. Damage to the nasal mucosa, the turbinates and the nasopharyngeal wall during nasotracheal intubation has been well described. Damage to the tube cuff is a frequent finding, especially when resistance is encountered during passage of the tube; this may necessitate reintubation over stiff plastics, gum bougies or woven stylets with no guarantee that the new cuff will not encounter a similar fate.

We have used a well lubricated Penrose drain, introduced into the nasal cavity, as a shield between the nasal structures and the advancing tube. An angulated forceps is used to insert the drain which is stabilized with two mosquito forceps. The endotracheal tube is inserted through the nasal cavity inside the Penrose drain (Figure).

FIGURE Endotracheal tube inserted inside Penrose drain.

After successful intubation, the Penrose drain is left *in situ* and is removed, together with the endotracheal tube, at the end of the surgical procedure.

We believe that the technique is beneficial in many patients, especially those with a difficult airway.

S. B. Backman MD PhD
M. Abou-Madi MD FRCPC
D. Trop MD FRCPC
Montreal Neurological Hospital
Montreal

Pain relief in labour

To the Editor:

"Giving an anaesthetic and not seeing the patient post-operatively is like doing an experiment without bothering to read results."* Yet, so few anaesthetists see their patients after anaesthesia. Small wonder that many of our patients complain despite our "best efforts."

Obstetric anaesthesia is a special situation. Pain in labour is one of the most severe forms of pain and, moreover, it occurs in those whom we love most.¹ We started an epidural service in 1957. Recently, we evaluated the efficacy of our technique in 65 parturients. All were healthy and at term. After a test dose of 2 ml, 2% lidocaine HCl, 10 ml 0.25% bupivacaine or 10 ml 0.25% bupivacaine were injected with 1 $\mu\text{g} \cdot \text{kg}^{-1}$ fentanyl. Demographic data were collected (Table) and the patient followed by an independent observer. Assessment was by Visual Analogue Scales (VAS) 20 min after epidural initiation, and then every hour until delivery, and again between 12 to 24 hr post delivery. Our results are presented in the Figure. The group of patients receiving bupivacaine and fentanyl mixture required fewer top-ups for the first two hours after the epidural was started.

Twelve to 24 hours after delivery we asked the patients to classify the quality of their pain relief during labour, during pushing, and during delivery as excellent, very good, good, fair and poor. Ninety-eight percent of women described their pain relief as good to excellent during labour, 74% as good to excellent during pushing and 94% as good to excellent during delivery.

Our audit confirmed the clinical impression that most patients were satisfied. We encourage other anaesthetists to do similar audits. It is the simplest and best way of

*Personal communication, James Parkhouse.