trials. These studies must be performed carefully and analyzed conservatively if one is to avoid making premature, incorrect conclusions. In particular, it will be important to resist the temptation to use this type of analysis for comparing the "quality" of anaesthesia among different hospitals, unless one can control for *all* the institution-specific factors which may influence patient outcome.

Peter H. Norman
M.D. Daley
Department of Anaesthesia
Toronto General Hospital
University of Toronto

REFERENCES

- 1 Cohen MM, Duncan PG, Tweed WA, et al. The Canadian four-centre study of anaesthetic outcomes: I. Description of methods and populations. *Can J Anaesth* 1992; 39: 420-9.
- 2 Cohen MM, Duncan PG, Pope WDB, et al. The Canadian four-centre study of anaesthetic outcomes: II. Can outcomes be used to assess the quality of anaesthesia care? *Can J Anaesth* 1992; 39: 430-9.
- 3 Duncan PG, Cohen MM, Tweed WA, Biehl D, Pope WDB, Merchant RN. The Canadian four-centre study of anaesthetic outcomes: III. Are anaesthetic complications predictable in day surgical practice? *Can J Anaesth* 1992; 39: 440-8.
- 4 Glantz SM. *Primer of Biostatistics*. 2nd ed. New York: McGraw-Hill, 1987.

REPLY

We thank Drs. Norman and Daley for taking the time to write about our study. There are several points to which we wish to respond.

The first issue raised relates to patient interviews. We disagree that "patients are often unreliable historians." It is more likely that health professionals do not ask questions in a manner which will solicit the desired information. Questions tend not to be asked in a systematic fashion during day-to-day clinical practice and comments such as "is everything alright" are very vague. It appears to be true that prompting the patient elicits higher reporting of rates of discomfort symptoms. There are several reasons for this. First patients do not want to "bother" busy doctors and nurses with complaints, even sometimes if they are troublesome and, as the authors point out, patients often expect to undergo some unpleasantness when having an operation. If asked "do you (or did you) have any problems with your anaesthetic," patients may only remember the most severe or unexpected symptoms or may be too inhibited to mention problems. Alternatively they may be so pleased to have "made it through" that lesser complaints are ignored. General questions concerning comfort and well-being lack the sensitivity to detect morbidity and negative answers cannot ensure that a given outcome was absent. In order to determine the actual rate of discomfort symptoms (whether severe or not), asking

the patient is the best (and often the only) method. In improving the quality of anaesthesia care, a mild degree of discomfort should not be the goal - elimination of symptoms altogether should be the ultimate aim.

In our study we used a systematic approach to patient interviews. Every patient was asked questions in a very similar manner and patients who could not understand the questions or who were too ill were excluded from the analysis. Thus we do not think that the results we achieved in the third paper were due to factors alluded to by Drs. Norman and Daley. However, we do note in the paper that the response rate to the interviews was very low and non-representativeness of the patients may have given rise to certain results. For example it may be that only the very sick patients were at home to receive phone calls the day after surgery thus explaining the high rate of symptoms in this population.

Drs. Norman and Daley also allude to the difficulty in performing studies of this type (specifically relating to reliability and validity) and point out that only "very preliminary conclusions" should be derived. We could not agree more. Although not necessary, we elected for various reasons to publish a separate paper on the methodological issues arising from such a study. First, we wanted to identify for anaesthetists how to perform this type of study, since "outcome" studies are rare in the field. Second, the conclusions of similar studies done in virtually every department (under the rubric of "audit" or "QA") are often accepted without challenge, but lack validity for the reasons we discuss. Finally, by describing our methods in some detail, we challenge other researchers to design studies which overcome some of these difficulties. Just as anaesthetists are not using the same drugs and protocols as they did even five years ago, clinical epidemiologists continuously update their study methods. In many large, multisite studies, the researchers have not described their protocols in nearly the detail which we did. This does mean that they did not confront similar issues but rather only that they went unrecognized.

With respect to the low kappa value across physician case reviewers, we agree that the value was less than desirable. We did not evaluate the reliability of the "causation assessment for real cases" because we used a consensus process. This was for pragmatic reasons: it was virtually impossible to get 12 busy clinician volunteers to review all cases, and we did not have adequate funds for training sessions for the reviewers. Future studies will have to pay case reviewers for their time and training. This was done in two studies quoted in the paper and would likely account for their high interrater reliability. However, despite the low kappa values we decided to use case reviewers to reflect actual clinical practice. Reviewers for morbidity/mortality rounds are rarely trained and usually only one person performs this function for each case. Yet in clinical practice, the "reliability" of this opinion is rarely questioned.

The next issue relates to the P values used for setting statistical significance in this study. While we agree with the authors, we wish to point out that the P values reported in the tables were those factors considered statistically significant at the P less than or equal to 0.01. The reader might infer that all the values reported were significant only at the 0.01 level, but that is the minimum which we accepted. Most of the values were much lower than this (see examples in the Table which correspond to Table VI in the third paper).¹ Not all statistical experts agree with the use of the Bonferroni correction for all situations. The correction tends to be too conservative if the number of comparisons is very high (i.e., no relationships

TABLE Adjusted relative odds for any intraoperative complications by preoperative medical conditions

Condition	Adjusted relative odds	P value
Asthma	2.2	0.0007
COPD	2.2	0.0005
Other respiratory	1.9	0.0004
MI < 6 months	6.4	0.03
MI > 6 months	1.1	0.76
Hypertension	2.2	0.0001
Arrhythmia	2.0	0.07
CHF	2.7	0.06
Other cardiovascular	1.2	0.53
Cerebrovascular	1.2	0.82
Gastrointestinal	2.7	0.0001
Renal	0.65	0.56

would be "statistically significant"). We would suggest, instead, that future papers (not only in anaesthesia, but in all journal articles) should report actual P values whenever possible and the reader can decide whether the association is meaningful or not.

Finally, we wish to make a few points about studies which employ non-randomized designs. First, randomized designs cannot be used to study "outcomes" of actual practice but can only be used to study specific research questions where groups of patients can be randomized. In attributing causal relationships one cannot use the results of only one study.² The reader must consider the results of similar studies and look for similar trends and results. Only after a number of studies have been performed can the reader comfortably assume a true relationship between a variable or interest and a particular outcome. Thus we do not agree with the authors that one should ignore relationships and associations we have described, but to evaluate each on its own merits and compare the results with other such studies. Progress in science is seldom made when accepted observations are limited to those we already understand.

Having said this we do agree with the authors that this type of analysis is not definitive for comparing the quality of anaesthesia across hospitals. We confirmed that institutions are different, and that traditional methods used to assess outcome of care are not applicable to anaesthesia. We argue that other methods must be developed for the specialty.

How then should the three studies we reported be used? First, they should be used to inspire the advancement of methodology of assessing "quality" in anaesthesia. Second, individual rates of adverse outcomes may be interesting in themselves and prompt a critical examination of different practices that account for rate discrepancies. Third, risk factor profiles derived by such studies should allow practising clinicians to focus on those patients in their daily practise.

Marsha M. Cohen MD FRCPC
Clinical Epidemiology Unit
Sunnybrook Health Science Centre
and the Department of Health Administration
University of Toronto, Toronto, Ontario
Peter G. Duncan MD FRCPC
Department of Anaesthesia

Royal University Hospital
Saskatoon, Saskatchewan

REFERENCES

- 1 Duncan PG, Cohen MM, Tweed WA, et al. The Canadian four-centre study of anaesthetic outcomes: III. Are anaesthetic complications predictable in day surgical practice? Can J Anaesth 1992; 39: 440-8.
- 2 Hill AB. The environment and disease: association or causation? Proc Roy Soc Med 1965; 58: 295-300.

Cost of soda lime

To the Editor:

We found Dr. Dion's letter on calculating the cost of inhalational anaesthesia¹ very interesting. According to his formulae, for a given percentage of vapour and for a given duration of anaesthesia, the cost varies directly with the fresh gas flow. This does not take into account the cost of consumption of soda lime at low fresh gas flows where rebreathing occurs. We would like to add to Dr. Dion's formulae accordingly. To do so the following additional information is required.

Cost (Sodasorb)	48.46¢ / 100 g Sodasorb
CO ₂ absorption capacity (Sodasorb)	45 g CO ₂ / 100 g Sodasorb ²

Using the molar volume of gas at 21° C of 24.12 L · mole⁻¹ and the molecular weight of CO₂ of 44.009 g · mole⁻¹, we calculate the cost of soda lime for complete rebreathing as follows:

$$CS \text{ (cents)} = 1.965 \dot{V}CO_2 T$$

where $\dot{V}CO_2$ is the CO₂ production in L · min⁻¹ and T is the duration of anaesthesia in minutes. The fraction of this that is rebreathed depends on the minute ventilation ($\dot{V}$) and the fresh gas flow (F), both in L · min⁻¹. For $F \geq \dot{V}$ one can assume that there is no rebreathing³ and obtain:

$$\text{fraction of expired gas rebreathed} = 1 - \frac{F}{\dot{V}}$$

The cost of soda lime as a function of flow is therefore;

$$\begin{aligned} CS \text{ (cents)} &= 1.965 \dot{V}CO_2 T \left(1 - \frac{F}{\dot{V}} \right) \\ &= -1.965 \left(\frac{\dot{V}CO_2}{\dot{V}} \right) TF + 1.965 \dot{V}CO_2 T \end{aligned}$$

Note that for $F \geq \dot{V}$, no soda lime is needed, so CS = 0.