REFERENCES

- 1 Atkinson RS, Rushman GB, Lee JA. A Synopsis of Anaesthesia. 8th. ed. Bristol: John Wright & Sons. 1977.
- 2 Miller RD. Anesthesia. 2nd. ed. New York: Churchill Livingstone. 1986.
- 3 Grey TC, Nunn JF, Utting JE. General Anaesthesia. 4th. ed. London: Butterworths. 1980.
- 4 Churchill-Davidson HC. A Practice of Anaesthesia. 5th. ed. Chicago: Year Book Medical Publishers Inc. 1984.
- 5 Collins VJ. Principles of Anesthesiology. 2nd ed. Philadelphia: Lee & Febiger 1976.
- 6 Badenoch AW. Manual of Urology. 2nd. ed. New York: William Heinemann. 1974.
- 7 Blandy J. Lecture notes on Urology. 3rd. ed. Oxford: Blackwell 1982.
- 8 Campbell MF, Harrison JH. Urology. 3rd. ed. Philadelphia: W. B. Saunders. 1970.
- 9 Devine CJ, Stecker JF. Urology in Practice. Boston: Little, Brown. 1978.
- 10 Winter CC. Practical Urology. Saint Louis: C. V. Mosby. 1969.
- 11 Smith DR. General Urology. 10th ed. Los Altos, Calif. Lange Medical. 1981.
- 12 Bodner H. Diagnostic and Therapeutic Aids in Urology; 230. Springfield: Charles C. Thomas. 1974.
- 13 Benzon HT, Leventhal JB, Ovassapian A. Ketamine treatment of penile erection in the operating room. Anesth Analg 1983; 62: 457-8.
- 14 Soliman MG, Tremblay NA. Nerve block of penis for postoperative pain relief in children. Anesth Analg 1978; 57: 495-8.
- 15 Comarr AE. Sexual concepts in traumatic cord and cauda equina lesions. J Urol 1971; 106: 375-8.
- 16 Weiss HD. The physiology of penile erection. Ann Intern Med 1972; 76: 793-9.

Percutaneous transtracheal ventilation for laser endoscopic procedures in infants and small children

To the Editor:

I believe that the method suggested by Ravussin and his colleagues in their recent article¹ may be extremely hazardous.

I cannot accept that on the basis of a report of only two cases a technique can be labelled as "... safe and useful ...". Some years ago Dr Fearon and I drew attention to the dangers of intratracheal catheter jet ventilation in children with obstructive airway lesions.² Since that time I have been made aware of further deaths resulting from the use of this method. To these dangers the present authors have

added those of percutaneous tracheal puncture. They are also advocating the use of a jet distal to a major obstruction lesion, a practice which is generally thought to be contraindicated.³ True they have added the use of an "exit route" for insufflated gases, and they stress vigilance and "fail safe" systems to minimise the risk of barotrauma. I am concerned, however, that such precautions are not infallible and that their method introduces some unnecessary potential dangers when much safer and simpler alternative methods and anaesthesia are available.

David J. Steward, MB FRCP
Department of Anaesthesia
British Columbia's Children's Hospital
Vancouver, British Columbia
V6H 3V4

REFERENCES

- 1 Ravussin P, Bayer-Berger M, Monnier P, Savary M, Freeman J. Percutaneous transtracheal ventilation for laser endoscopic procedures in infants and small children with laryngeal obstruction: Report of two cases. Can J Anaesth 1987; 34: 83-6.
- 2 Steward DJ, Fearon B. Anaesthesia for laryngoscopy, Br J Anaesth 1981; 53: 320.
- 3 France NK. Anaesthesia for pediatric ENT. In: Pediatric Anesthesia, Gregory G.A. (Ed). Churchill Livingstone, New York, 1983.

REPLY

We appreciate Dr Steward's comments on our recent article, and agree with his concern about the possible complications using percutaneous transtracheal jet ventilation. However, we believe that adequate precautions were undertaken to minimise these risks, and that there is a significant added advantage in avoiding tracheostomy in small children. Furthermore, a catheter introduced into the trachea under endoscopic control allows at any time simple insufflation of oxygen ($1-3 \text{ L} \cdot \text{min}^{-1}$) should complete obstruction occur. We wish to emphasise that we do not suggest this technique for routine cases where the use of a more conventional anaesthetic technique is possible, but that it should be reserved for use in specific cases such as described. Laser surgery, unlike endoscopy, requires good surgical access to an immobile operative site, usually achieved with the aid of muscle relaxation. However, the importance of an "exit route" for the insufflated gases can not be over-emphasised,¹⁻³ and a bronchoscope in place is a perfect guarantee. Thus in certain rare cases we have found the technique we described to be a valid alternative in experienced hands. We would be interested to know the "safer and simpler alternative methods of anesthesia" that Dr Steward would have used in the cases reported, in particular case I, the 4-month-old infant with a 25 mm long subglottic haemangioma obstructing the tracheal lumen by approximately 80 per cent while at the same ensuring surgical access to an immobile operating field.

P. Ravussin MD FMH
Agrégé
CHUV, 1011 Lausanne, Switzerland

REFERENCES

- 1 Carden E, Becker G, Hamood H. Percutaneous jet ventilation. *Ann Otol* 1976; 85: 652-5.
- 2 Smith RB, Myers EN, Sherman H. Transtracheal ventilation in paediatric patients. *Br J Anaesth* 1974; 46: 313-4.
- 3 France NK. Anaesthesia for Pediatric ENT. In: Pediatric Anesthesia, Gregory GA (Ed). Churchill Livingstone, New York, 1983.

Epinephrine test dose for confirmation of intravascular migration of epidural catheter in a parturient

To the Editor:

We would like to report a case illustrating the probable intravascular migration of an epidural catheter which presented as tinnitus during injections and was confirmed by use of an epinephrine test dose via the catheter.

A 22-year-old primiparous term parturient requested epidural analgesia after eight hours of labour and at five cm cervical dilatation. A Tuohy epidural needle was inserted at the L₄₋₅ interspace and 4 ml of 0.5 per cent bupivacaine was injected via the needle. A catheter (Portex, 18G nylon catheter, 3 eyes) was inserted and an additional 4 ml of 0.5 per cent bupivacaine was injected via the catheter. The patient had no systemic complaints during the injection and had good analgesia for two hours, after which she requested a second injection. Five ml of 0.5 per cent bupivacaine was given. The parturient immediately complained of ringing in the ears which disappeared after a few minutes. After no improvement in pain relief in 30 minutes an additional injection of only 2 ml of 0.5 per cent bupivacaine again caused ringing in the ears.

Aspiration of the epidural catheter did not yield any blood. Because of our suspicion of an inadvertent intravenous injection, 10 µg of epinephrine were injected into the epidural catheter during ECG monitoring. A uterine contraction began soon after the injection and so the increased heart rate could not be properly evaluated. After four minutes, an additional 10 µg of epinephrine were injected into the epidural catheter with a subsequent increase in heart rate from 66 to 86 beats·min⁻¹ after 30 seconds. After another four minutes' wait, a third injection of 10 µg of epinephrine into the epidural catheter caused the heart rate to increase from 74 to 88 beats·min⁻¹ after 26 seconds. Systolic blood pressure increased from 110 to 140 mmHg over the course of the three injections. Finally, a 5 ml saline placebo injection produced neither tinnitus nor an increase in heart rate.

The epidural catheter was removed and on inspection only the proximal hole was patent, with the distal two holes being plugged with blood. A second epidural catheter placed at the L₂₋₃ interspace provided excellent analgesia with no systemic complaints.

The addition of epinephrine to an epidural test dose was described by Moore and Batra¹ who showed that an intravenous injection of 15 µg of epinephrine caused an increase in heart rate of about 30 beats·min⁻¹ within an average of 30 seconds after injection. In our case, we injected only 10 µg of epinephrine, although we repeated the injection three times. The heart rate increased 20 beats·min⁻¹ within 30 seconds of the second injection and 14 beats·min⁻¹ within 26 seconds of the third injection. These responses clearly suggest that the epinephrine was injected intravascularly via the epidural catheter.

Abraham *et al*² reported on eight parturients out of 250 who exhibited an increase in heart rate following the injection of a lidocaine, epinephrine and dextrose containing test dose via an epidural catheter. They presumed those to be accidental intravenous injections.

Cartwright *et al*³ subsequently showed however that the epidural injection of bupivacaine alone will cause an increase in maternal heart rate in 12 per cent of patients. Our patient received epinephrine alone without any local anaesthetic and showed no response to a placebo saline injection.

Therefore, when testing for the accidental intravenous placement of an epidural catheter we would suggest using epinephrine alone. The ideal single epidural test dose that will clearly demonstrate both intravenous and subarachnoid injections remains to be established.

Joseph Eldor MD

David Z.N. Frankel MD FRCP

David Olshawang MD

Department of Anesthesia

Shaare Zedek Medical Center

Jerusalem, Israel

REFERENCES

- 1 Moore DC, Batra MS. The components of an effective test dose prior to epidural block. *Anesthesiology* 1981; 55: 693-6.
- 2 Abraham RA, Harris AP, Maxwell LG, Kaplow S. The efficacy of 1.5 per cent lidocaine with 7.5 per cent dextrose and epinephrine as an epidural test dose for obstetrics. *Anesthesiology* 1986; 64: 116-9.
- 3 Cartwright PD, McCarroll SM, Antzaka C. Maternal heart rate changes with a plain epidural test dose. *Anesthesiology* 1986; 65: 226-8.