by the authors' positive evaluation of this simulator system.

Two other items should be called to the attention of the reader. First, I find it puzzling that the authors did not mention that a simulation centre using a simulator from CAE-Link (the company is now known as CAE Electronics Incorporated) had been under design and construction at Sunnybrook Health Science Centre/University of Toronto since the summer of 1994. This facility opened several months ago and is currently engaged in a variety of training and research activities using the simulator, including the intensive training course in Anaesthesia Crisis Resource Management. Second, in their discussion of the use of simulation for training in anesthetic crises I believe that mention should be made of the special simulator-based curriculum on Anesthesia Crisis Resource Management (ACRM)¹⁻³ that was pioneered by my laboratory in 1990, and that is currently being offered at three simulator centers affiliated with major teaching institutions^{1,2,4,5} in North America (Stanford University, Harvard University, University of Toronto).

David M. Gaba MD
Stanford University School of Medicine
Palo Alto Veterans Affairs Medical Center

REFERENCES

- 1 Howard SK, Gaba DM, Fish KJ, Yang GS, Sarnquist FH. Anesthesia crisis resource management training: teaching anesthesiologists to handle critical incidents. *Aviat Space Environ Med* 1992; 63: 763-70.
- 2 Holzman RS, Cooper JB, Gaba DM, Philip JH, Small S, Feinstein D. Participant responses to realistic simulation training in anesthesia crisis resource management (ACRM). *Anesthesiology* 1993; 79: A1112.
- 3 Gaba DM, Botney R, Howard SK, Fish KJ, Flanagan B. Interrater reliability of performance assessment tools for the management of simulated anesthetic crises. *Anesthesiology* 1994; 81: A1277.
- 4 Holzman RS, Cooper JB, Gaba DM, Philip JH, Small S, Feinstein D. Anesthesia crisis resource management: real-life simulation training in operating room crises. *J Clin Anesth* (in press).
- 5 Kurrek MM, Small S, Feinstein D, Russell R, Fish K, Gaba D. Anaesthesia Crisis Resource Management (ACRM) instructor course - teaching the teachers (abstract). Presented at the Conference on Simulators in Anesthesia Education, Rochester, NY, 1995.

REPLY (2)

A check of our final submitted manuscript revealed that we had originally written "The simulation technology used in the system was acquired in prototype form from Dr. Gaba and

Dr. Williams of Stanford University, with contributions (mostly pharmacokinetic and pharmacodynamic models) from Dr. Schwid of the University of Washington." The change to "We acquired a prototype ..." was later made at the CJA editorial office and unfortunately escaped our notice. As stated, our assessment was based on using the unit on a trip to the CAE factory in Binghamton, New York.

Dr. Gaba was puzzled by our failure to mention the simulator at the Sunnybrook Health Science Centre. However, at the time our original manuscript was submitted, Sunnybrook had not even been chosen as a simulator site. Indeed, the main purpose of our assessment of the CAE system was to explore the possibility of acquiring it for the Toronto Hospital. We are, however, very pleased that Sunnybrook has since taken on a strong leadership role in providing anaesthesia simulator services to Canadians.

Finally, we regret that tight space limitations did not allow us to cite the abstracts on Anaesthesia Crisis Resource Management to which Dr. Gaba refers. We are sure that when they are eventually published as full papers, they will be valuable contributions to the anaesthesia literature.

D. John Doyle MD PhD FRCPC
Ramiro Arellano MD FRCPC
Department of Anaesthesia
The Toronto Hospital

Respiratory depression associated with patient-controlled analgesia

To the Editor:

Respiratory depression associated with patient-controlled analgesia (PCA) was recently reviewed¹ with an accompanying editorial.² Risk factors included concurrent use of background infusions, advanced age, concomitant administration of sedative/hypnotic medications, pre-existing sleep apnoea syndrome, and excessively liberal prescribing. We report a patient who suffered severe respiratory depression when epidural analgesia was changed to *iv* PCA.

The patient, a 44-yr-old, ASA 2, 74 kg man underwent partial gastrectomy for gastric carcinoma. Before general anaesthesia, an epidural catheter was inserted at T₉₋₁₀. Intraoperative analgesia was provided with epidural bupivacaine (76.25 mg) and epidural fentanyl (150 µg). Post-operative analgesia was with a continuous epidural infusion of bupivacaine 0.1% with meperidine 2 mg · ml⁻¹ adjusted, after initial titration, to 7.5 ml · hr⁻¹. After surgery, he was pain-free and satisfied. The following morning, whilst comfortable, the epidural catheter was removed for technical reasons. Further analgesia was with *iv* morphine by PCA (bolus 2 mg, lockout 5 min, four-hour maximum 30 mg) and no background infusion. Seven hours later, the patient was found unresponsive