REFERENCES

- 1 Butscher K, Mazoit JX, Samii K. Can immediate opioid requirements in the post-anaesthesia care unit be used to determine analgesic requirements on the ward? *Can J Anaesth* 1995; 42: 461-6.
- 2 Moote CA. Postoperative pain management – back to basics (Editorial). *Can J Anaesth* 1995; 42: 453-7.

REPLY

Going "beyond basics" in health care is an issue which is currently undergoing intense debate. It is time to define core and comprehensive services.¹ Which services are necessary and which are a luxury? Which services should the public be asked to fund and for which services should the patient be required to pay? The cornerstone for such technology assessment must be evidence-based medicine. New therapeutic procedures must be compared with standard treatments considering safety, efficacy and cost.

Safety – Respiratory arrest² and severe respiratory depression³ can and does occur with PCA. Problems have occurred when well-meaning nurses, family and visitors help the patient push the button. Equipment problems are an additional concern. An ampoule of naloxone is routinely at the bedside for patients using PCA. This has never been standard practice for im analgesia.

Efficacy – Changing the technique of administration does not dramatically alter the efficacy or safety of morphine. Papers that compared im and PCA analgesia, have found both to be equally effective.⁴ Inadequate analgesia has also been reported with PCA.⁵

Cost – The Committee on Pain Management under the Section on Clinical Care in the American Society of Anesthesiology has developed pain outcome measures for pain management. In the list of nine pain outcome measurement categories the first category listed is cost.⁶ If conventional nurse administered opioid analgesia produces a similar outcome at lower cost, it should be adopted as the analgesic technique of first choice. In a health care system funded solely by tax dollars, PCA should be reserved for patients who cannot be managed by nurse administered analgesia.

Carol A. Moote, MD FRCP
London, Ontario

REFERENCES

- 1 Wyman M, Feeley J, Brimacombe G, Doucette K. Core and comprehensive health care services: 4. Economic issues. *Can Med Assoc J* 1995; 152: 1601-4.
- 2 Zimmerman DL, Stewart J. Postoperative pain management and Acute Pain Services activity in Canada. *Can J Anaesth* 1993; 40: 568-75.
- 3 Etches RC. Respiratory depression associated with patient-controlled analgesia: a review of eight cases. *Can J Anaesth* 1994; 41: 125-32.
- 4 Wheatley RG, Shepherd D, Jackson IJ, Madej TH, Hunter D. Hypoxaemia and pain relief after upper abdominal surgery: comparison of i.m. and patient-controlled analgesia. *Br J Anaesth* 1992; 69: 558-61.
- 5 Smythe MA, Martin SE, Schad RF. Retrospective evaluation of patient-controlled analgesia use in a large teaching institution. *DICP* 1991; 25: 1058-61.
- 6 Thompson GE. Outcome measures defined for pain management. *American Society of Anesthesiologists Newsletter* 1995; 59: 31-2.

Medication labels: for whose benefit?

To the Editor:

We wish to report a "near-miss" drug error that was averted because of the report by Ahdal and Bevan.¹ These authors described a case of prolonged neuromuscular blockade following an overdose of clindamycin. Misinterpretation of the drug label caused the overdose and contributed to a similar incident described below.

Recently, one of the authors (R.C.) was asked to administer clindamycin (300 mg) to a patient undergoing general anaesthesia. The prominent text on the clindamycin vial, printed in large blue letters, was "Dalacin C Phosphate." The number "150" was the second most prominent feature. In smaller font, "clindamycin phosphate injection USP" was evident. The contents of the two vials (8 mL) were drawn into a syringe in order to prepare a dose of 300 mg. The need to open two vials seemed unusual and the report of Ahdal and Bevan came to mind. Upon closer inspection of the vials, "clindamycin/mL" was noted in small print. The syringe, therefore, contained 1200 mg, four times the required dose. The dose of clindamycin reported by Ahdal and Bevan was also increased by a factor of four, suggesting Dalacin C Phosphate was involved.

This label (Figure) meets the standards set by the Canadian Society of Hospital Pharmacists, yet it is difficult to read.

FIGURE