REFERENCES

- 1 *Morris IR*. Fiberoptic intubation. *Can J Anaesth* 1994; 41: 996–1007.
- 2 *Cold AFD, Mallon JS, Rolbin SH*. Fiberoptic intubation [Letter]. *Can J Anaesth* 1995; 42: 839–40.
- 3 *Benumof JL*. Management of the difficult adult airway. With special emphasis on awake tracheal intubation. *Anesthesiology* 1991; 75: 1087–110.
- 4 *Cormack RS, Lehane J*. Difficult tracheal intubation in obstetrics. *Anaesthesia* 1984; 39: 1105–11.
- 5 *Ovassapian A, Mesnick PS*. The art of fiberoptic intubation. In: Sandler AN, Doyle DJ (Eds.). *Anesthesiology Clinics of North America*. Philadelphia: WB Saunders, 1995; 39: 1–409.

Errata

Maltby JR, Loken RG, Beriault MT, Archer DP. Laryngeal mask airway with mouth opening less than 20 mm. *Can J Anaesth* 1995; 42: 1140–2.

Please note that the last sentence of the “Discussion” on page 1142 should have read:

We conclude that the LMA may be considered in patients with restricted mouth opening for guided intubation or for definitive airway *management during* surgery and during recovery from anaesthesia.

Brimacombe J. The advantages of the LMA over the tracheal tube or facemask: a meta-analysis. *Can J Anaesth* 1995; 42: 1017–23.

Please note that the last sentence (bottom page 1020, first column) should have read:

Surprisingly the LMA *does not* cause an increased work of breathing compared with the TT even though the larynx, which contributes 25% of total airways resistance, is not bypassed.⁷⁰