humain brain slices by histochemical methods.³ Based on these findings, and with the inadequate level of current knowledge on the actual number of active enzyme receptors and substrate concentrations in vivo at distinct cholinergic receptor sites, one cannot rule out the practical clinical value of this "academic" finding of a moderate PChE inhibitory effect.

Elemer K. Zsigmond, MD
Department of Anesthesia
University of Illinois
College of Medicine at Chicago, Chicago, Illinois

REFERENCES

- 1 Foldes FF, Smith JC. The inhibition of human cholinesterases with anticholinesterases used in the therapy of myasthenia gravis. *Ann NY Acad Sci* 1966, 135: 287.
- 2 Foldes FF, Molloy RE, Zsigmond EK, Zwartz JA. Hexafluorenum: its anticholinesterase and neuromuscular activity. *J Pharmacol Exp Ther*, 1960, 129: 400.
- 3 Radic Z. Effect of atropine on esterases of human blood and pig liver. Abstract #4 p. 43, VI. Balkan Biochemical and Biophysical Days, Plovdiv, April 15-19, 1985. Bulgaria.
- 4 Koelle WA, Koelle GB. The localization of external or functional acetylcholinesterase at the synapses in autonomic ganglia. *Progr Brain Res* 1971, 34, 371.
- 5 Foldes FF, Zsigmond EK, Foldes VM. The distribution of acetylcholinesterase and butyrylcholinesterase in the human brain. *J Neurochem* 1962, 9: 559-72.

Nuffield Department of Anaesthetics, Oxford

To the Editor:

A scientific meeting will be held in Oxford from 23rd-25th July, 1987, to celebrate the Fiftieth Anniversary of the founding of the Nuffield Department of Anaesthetics. This meeting will follow the Second World Congress on the History of Anaesthesia, which is to be held in London from 20th-23rd July, 1987.

Anaesthetists and others who have had connections with the department are invited to write to:

Nuffield Department of Anaesthetics (Dept. GP/87),
Radcliffe Infirmary,
Woodstock Road,
Oxford,
OX2 6HE,
England

enclosing their full name and postal address so that they may be circulated with further particulars.

Professor M.K. Sykes
University of Oxford

Transtacheal ventilation

To the Editor:

I was interested to read the report by Ravussin and Freeman on transtracheal ventilation.¹ Two points are worth raising, firstly that high frequency ventilation is not necessary to produce oxygenation. A simple hand-operated trigger device such as the Sanders bronchoscope injector can be used in most instances equally well^{2,3} and is more likely to be available in the emergency situation.

Secondly, in the interests of safety, I would like to stress the importance of a clear upper airway to allow the inflated gases to escape. This is best achieved with the patient's neck in the extended position (as for thyroidectomy). In my experience (in 60 cases) the surgeon can easily flex the head and neck while working on the lower jaw for example, resulting in obstruction of the gas escape pathway. Extra vigilance is necessary when using transtracheal ventilation, to prevent the occurrence of high pressure in the lower airways, leading to possible pneumothorax and emphysema.

P.R. Layman MB CHB FFARCS
Department of Anaesthetics
John Radcliffe Hospital
Headington, Oxford

REFERENCES

- 1 Ravussin P, Freeman J. A new catheter for transtracheal ventilation and resuscitation. *Can Anaesth Soc J* 1985; 32: 60-4.
- 2 Layman PR. Transtracheal ventilation in oral surgery. *Annals of the Royal College of Surgeons* 1983; 65: 318-20.
- 3 Layman PR. Bypassing a problem airway. *Anaesthesia* 1983; 38: 478-80.

REPLY

We thank Dr. Layman for his letter. He points out two major aspects of great importance: (1) that when transtracheal jet ventilation is used, a clear upper airway is

mandatory, because any gas injected into the trachea must be able to escape. This aspect is stressed in the discussion of our paper but it cannot be over-emphasized. (2) Transtracheal ventilation through a 13 G or 14 G catheter inserted into the trachea must be a jet-type ventilation. I entirely agree that high frequency jet ventilation (HFJV) is not the only way to achieve good gas exchange. The Sanders Jetting Device¹ or the Carden Jetting Device² are both cheap, hand-operated and not cumbersome. Both devices can be plugged into a compressed oxygen supply system. I personally prefer the Carden Jetting Device because it has a pressure regulator (as in the HFJV ventilators) and thus the 405 kPa (4 bar) pressure from the oxygen wall gas outlet can be adjusted according to the size and the condition of the patient.

The catheter described in our paper was first designed for easy, rapid use in an emergency but increasingly we use it for endoscopic and laser procedures involving the larynx or the very upper part of the trachea. Oedema often develops around the glottis after such procedures, and the catheter should be left in place for up to 12 hours, in case an obstruction of the glottis should develop. The catheter is very well tolerated by the patients even when they are fully recovered and conscious after anaesthesia.

P. Ravussin MD FMH
Department of Anaesthesia
Centre Hospitalier Universitaire Vaudois
Lausanne/Switzerland

REFERENCES

- 1 Sanders RD. Two ventilating attachments for bronchoscopes. *Del Med J* 1967; 39: 170–75.
- 2 Carden E, West HR. Further advances in anesthetic techniques for microlaryngeal surgery. *Anesth Analg* 1974; 53: 584–7.