provide pre-induction data for the individual treatment groups.

Charles L. MacAdams MD FRCPC
Ian R. Thomson MD FRCPC
Department of Anesthesia
St. Boniface General Hospital
University of Manitoba
Winnipeg, Manitoba R2H 2A6

REFERENCE

- 1 Miller DR, Wellwood M, Teasdale SJ *et al.* Effects of anaesthetic induction on myocardial function and metabolism: a comparison of fentanyl, sufentanil and alfentanil. *Can J Anaes* 1988; 35: 219–33.

REPLY

We are grateful for the interest in our study expressed by Drs. MacAdams and Thomson. They carefully reviewed our manuscript and correctly identified limitations which were encountered in our assessment of myocardial contractility in patients undergoing narcotic anaesthesia. We concur with some of their observations. Volume loading before and after induction produced only a small increase in systolic blood pressure and thus limited our ability to assess systolic function. We also agree that the imprecision inherent in the method of determination of the left ventricular end-systolic volume index may have decreased our ability to assess systolic function. However, despite these considerations, we believe that there is compelling evidence for a decrease in myocardial contractility associated with narcotic anaesthesia.

First, the large variability associated with our measurements decreased the likelihood of our finding a statistically significant depression in systolic function. The fact that we were able to demonstrate such a difference suggests that the depression was profound. We were able to demonstrate a significant difference because we employed a powerful statistical tool. Analysis of covariance permitted the simultaneous evaluation of the slope and the position of the systolic pressure–volume relation. The figure in the letter from Drs. MacAdams and Thomson demonstrated that the systolic pressure–volume points post-induction were below the pre-induction values. This difference in position may represent a decrease in end-systolic elastance. Our sensitive method of analysis which incorporates both slope and position permitted us to conclude that this represented depression in systolic function.

Secondly, we believe that there is additional evidence for depressed myocardial contractility with narcotic anaesthesia induction. Preload recruitable stroke-work index also decreased with anaesthetic induction. Glower¹ and colleagues have demonstrated that this relation provides a better estimate of myocardial contractility than does end-systolic elastance. There are in vivo measurements, and we do not find it surprising that loss of consciousness with narcotic induction, and the accompanying loss of sensory input, is associated with a reduction in myocardial contractility. Such a demonstration is dependent upon the sensitivity of the measurements. The importance is in the comparisons. In an effort to increase the accuracy of these comparisons we have recently developed

on-line pressure-volume loop capability to pin-point more precisely ventricular end systole for future studies.

Finally, although the negative effects on cardiac function were observed, to varying degrees, with all three narcotics, the adverse effects on myocardial metabolism (and the greatest decrease in diastolic compliance) were observed only after alfentanil. The fact that all three groups were ventilated with equal positive pressure yet showed different diastolic compliance slopes suggests ischaemic-induced changes. This was supported by adverse metabolic changes in the alfentanil group.

In conclusion, we believe that anaesthetic induction with the narcotic agents studied produced small but significant alterations in myocardial function and metabolism (the latter in the alfentanil group) in patients undergoing elective coronary bypass surgery. Patients with impaired ventricular function and those who have acute ischaemia may have a more pronounced response to narcotic induction. Caution should be exercised under these circumstances.

Sallie Teasdale MD FRCPC
Donald Miller MD FRCPC
Richard Weisel MD FRCSC
Departments of Anaesthesia and Cardiovascular Surgery
Toronto General Hospital
University of Toronto
Toronto, Ontario

REFERENCES

- 1 Glower DD, Spratt JA, Snow ND *et al.* Linearity of the Frank-Starling relationship in the intact heart: The concept of preload recruitable stroke work. *Circ* 1985; 71: 994–1009.

Fentanyl, sufentanil, alfentanil and myocardial function

To the Editor:

Although it is stated in the article "Effects of anaesthetic induction on myocardial function and metabolism: a comparison of fentanyl, sufentanil and alfentanil" (*Can J Anaesth* 1988; 35: 219–33) that the patients in the alfentanil group had a progressive decrease in blood pressure for nearly five minutes, this is not shown in Figure 2. The mean arterial pressures in the alfentanil and sufentanil groups were equally depressed (compared with fentanyl) one minute after induction, but it is the sufentanil group which subsequently shows the persistently lowest blood pressures.

It is also stated that induction caused a reduction in cardiac index with all drugs but unfortunately this was not represented diagrammatically. Ejection fraction is shown, and this demonstrates the greatest increase in the alfentanil group. This cannot have been due to a reduction in afterload (systemic vascular resistance was not measured but mean arterial pressure according to Figure 2