Correspondence

NSAIDs

To the Editor:

I was interested to read the editorial by Code¹ on the use of NSAIDs in postoperative pain but I believe the effects of these drugs on renal function in the perioperative period was understated. The reference from Goodman and Gilman reflects primarily the general medical experience with these drugs when renal dysfunction is relatively rare except in the disease states listed where renal perfusion is dependent on the vasodilatory actions of prostaglandins. The situation in patients undergoing major surgery is different.

However healthy are our patients preoperatively and however well they are managed intraoperatively, many will be subjected to the neuro-humoral stress response to surgery involving the release of catecholamines, renin, angiotension and vasopressin (anti-diuretic hormone) which attain levels similar to those found in morbid disease states such as congestive heart disease or cirrhosis of the liver. Under normal circumstances these vasoconstrictor hormones stimulate the release of vasodilatory prostaglandins from the vascular endothelium of the afferent renal arteries and thus protect the kidneys from ischaemic injury. However, NSAIDs block the synthesis of these compounds² in patients undergoing oesophagogastrectomy. The importance of this effect was further demonstrated by Perttunen *et al*³ who compared diclofenac infusion with placebo in fluid restricted post-thoracotomy patients. Their control group passed a mean average of 230 ml of urine in the first 24 hr after surgery whereas the diclofenac group passed only 41 ml. Some consolation can perhaps be drawn from the finding in both of these studies that the effect was substantially less in the second 24 hr period suggesting that this is perhaps the time to be introducing this potent class of medication.

It is vital that we recognise the extent to which we stress the kidneys during and after major surgery and that we be circumspect in our decision to discard the protective mechanism that nature has deemed necessary to bestow upon these vulnerable organs.

Stephen F Jones MD
Specialist in Anaesthesia and Pain Management
Waikato Hospital
Hamilton, New Zealand

REFERENCES

- 1 Code W. NSAIDs and balanced analgesia. *Can J Anaesth* 1993; 40: 401-5.
- 2 Power I, Cumming AD, Pugh GC. Effect of diclofenac on renal function and prostacyclin generation after surgery. *Br J Anaesth* 1992; 69: 451-6.
- 3 Perttunen K, Kalso E, Heinonen J, Salo J. IV diclofenac in post-thoracotomy pain. *Anaesth* 1992; 68: 474-80.

REPLY

Thank you for your letter regarding my editorial "NSAIDs and balanced analgesia." It is important to outline a number of different interpretations which are possible from your letter. You wrote, "I believe the effects of these drugs on renal function in the perioperative period was understated." My editorial's goal was to enhance anaesthetists' use, or at least potential use, for NSAIDs in perioperative pain control. An editorial cannot be a review. As anaesthetists, we are fortunate that NSAIDs have been widely used for years. Data reveal that each year over 70 million prescriptions for NSAIDs are filled in the United States¹ and over 11 million in Australia.² Acute renal dysfunction associated with NSAIDs is usually reversible after discontinuation of the NSAID.³

If one considers the wealth of literature available on NSAIDs from disciplines outside of anaesthesia, then anaesthetists, as pharmacologists, have no excuse to be unaware of complications of NSAIDs. Unfortunately, there is minimal data about the short-term (less than one week) use of NSAIDs. What is available suggests they are safer in limited use than in prolonged use with respect to gastrointestinal bleeding.⁴ Renal function is likely to show minimal and clinically insignificant changes in healthy patients from short-term use.⁵

It is interesting that you quote the Perttunen article with thoracotomy and postoperative fluid restriction to suggest renal injury by diclofenac. Perioperative oliguria in the first 24 hr after major surgery is almost always induced by hypovolaemia. You noted the kidneys "recovered" in 24 hrs. Postoperative patients are now seldom treated with fluid restriction and intermittent boluses of furosemide for oliguria.

The paper, by Power *et al.*, examined the effect of NSAIDs on renal function after a very major procedure, oesophagogastrectomy, and measured prostaglandin concentrations directly related to renal function. The patients were elderly, had major surgery for a debilitating disease, and would be particularly susceptible to NSAID-induced renal toxicity. Power *et al.* noted that although diclofenac impaired renal function on the first day after surgery, this effect was short-lived.

In Canada and the United States, ibuprofen, acetylsalicylic acid (ASA) and acetaminophen are all available without prescription. Significant renal injury from use of these or prescribed NSAIDs for short-term use (two to three days) must be exceedingly rare in drugs used so frequently. Anaesthetists should be aware of all adverse effects of any drugs we use but should still use the safest combination for the best analgesic outcome. I still feel that NSAIDs should have an increasing role in pre-

vention and treatment of surgical pain. Patients with early renal failure NSAIDs are best avoided, or at least renal function should be carefully monitored.

William Code MD FRCP
Department of Anaesthesia
Cowichan District Hospital

REFERENCES

- 1 Murray MD, Brater DC. Renal toxicity of the nonsteroidal anti-inflammatory drugs. *Ann Rev Pharmacol Toxicol* 1993; 33: 435-65.
- 2 Day RO, Henry DA, Muirden KD, et al. Non-steroidal anti-inflammatory drugs induced upper gastrointestinal haemorrhage and bleeding. *Med J Aust* 1992; 157: 810-2.
- 3 Moote CA. Complication associated with NSAIDs use for acute pain management. *Perspectives in Pain Management* 1993; 3: 3-6.
- 4 Kehlet H, Dahl JB. Are perioperative nonsteroidal anti-inflammatory drugs ulcerogenic in the short term? *Drugs* 1992; 5: 38-41.
- 5 Aitken HA, Burns JW, McArdle CS, Kenny GN. Effects of ketorolac trometamol on renal function. *Br J Anaesth* 1992; 68: 481-5.

Tracheal tube leak test

To the Editor:

The paper on the leak test by Schwartz *et al.* was interesting. The conclusion that it is unreasonable to set an upper limit of leak pressure for changing all endotracheal tubes is contradicted by their results. Of the 242 patients studied 30 (12%) were excluded because one or both examiners found that no leak occurred at 50 cm H₂O. Does this mean the authors would consider a leak at 50 ($\pm$ 18) cm H₂O unacceptable?

Does the finding that the leak measured does not vary between observers invalidate the test? The need to determine fit of the endotracheal tube in the trachea and at the level of the cricoid ring derives from the fear that too tight a tube will cause pressure changes of the tissues at the level of the cricoid. These changes may cause transient problems such as stridor after extubation¹ or may be directly related to later tracheal stenosis.² The fact that a leak is detected at approximately 25 ($\pm$ 9.5) cm H₂O means that a pressure equal to the leak pressure is being exerted on the tracheal mucosa at the cricoid ring. This pressure should not exceed the perfusion pressure of the tissues at that level. In adults this has been estimated as 30 cm H₂O (22 mmHg).³ If we no longer measure the leak pressure, there is no other way we can easily estimate how tight the tube is. The leak pressure may increase appreciably⁴ in children that undergo cardiac surgery, this may be due to subglottic oedema as a reflection of generalised oedema. The increase in the

pressure may further compromise perfusion particularly if the arterial pressure is labile. A tube that fits well at the start of a case may become too tight. It is the duration and the age of the patient that seem to be the major determinants of tracheal stenosis. Thus the ex-premature neonate whose trachea has been intubated for some days is at particular risk.⁵ I know of no study of leak pressure in this population.

J.H. Smith MB MRCP FRCA
Newcastle-upon-Tyne
England

REFERENCES

- 1 Koka BV, Jeon IS, Andre JM, MacKay I, Smith RM. Post intubation croup in children. *Anesth Analg* 1977; 56: 501-5.
- 2 Minnigerode B, Richter HG. Pathophysiology of subglottic tracheal stenosis in childhood. *Prog Pediatr Surg* 1987; 21: 1-7.
- 3 Seegobin RD, van Hasselt GL. Endotracheal cuff pressure and tracheal mucosal blood flow: endoscopic study of effects of four large volume cuffs. *BMJ* 1984; 288: 965-8.
- 4 Smith JH, O'Kelly SW. Brain swelling after coronary artery surgery (Letter). *Lancet* 1993; 342: 1370-1.
- 5 Weber TR, Connors RH, Tracy TF Jr. Acquired tracheal stenosis in infants and children. *J Thorac Cardiovasc Surg* 1991; 102: 29-34.

REPLY

We thank Dr. Smith for his response to the paper entitled "Tracheal Tube Leak Test - Is There Inter-observer Agreement?" We stated that we believe it is unreasonable to set a rigid upper limit of leak pressure for the changing of all endotracheal tubes. We also suggested that other factors such as duration of surgery, difficulty of intubation, risk of aspiration of gastric contents, surgical position, and history of croup or other airway abnormalities be considered when deciding if tight-fitting endotracheal tubes should be changed.¹ The optimal limits of leak pressure used to avoid laryngeal injury are unknown. Koka *et al.* suggested using 25 cm H₂O, Litman and Keon use 40 cm H₂O.³ In this study it was found there is considerable variation between two experienced observers in assessing leak pressure. Rather than set a rigid upper limit, the peak test should be used as one of multiple factors employed to determine if the endotracheal tube should be changed. We agree with Dr. Smith that the plan to undergo cardiopulmonary bypass should be added to the list of factors used when deciding to change tight-fitting endotracheal tubes.

Dr. Smith questions why we excluded patients in which one or both examiners found no leak at 50 cm of water. We were reluctant to raise the intrathoracic pressure to greater levels for fear of barotrauma. Since no specific leak pressure was assigned by one or both of the examiners in these patients, no inter-observer difference could be calculated.

In summary, we did not intend to suggest that the leak pressure no longer be measured. We suggest that the absolute