Correspondence

Drug errors

To the Editor:

I was disappointed to read the recent clinical report by Orser and Oxorn.¹ This report would seem to prove that history is indeed condemned to repeat itself.

As a senior resident in Saskatchewan in 1984, I had the misfortune of a similar adverse event. In fact, I made the same error, the same way, with the same two drugs and ampoules while attempting to reverse residual neuromuscular blockade in a healthy woman. The accidental injection 1 mg epinephrine was followed by ventricular tachycardia and subsequent cardiac arrest. Like Orser and Oxorn, the patient was successfully resuscitated and made a full and complete recovery. Immediately after this adverse event, the ampoules used for neuromuscular reversal were examined. We were shocked to realize that one of the ampoules originally thought to be glycopyrrolate was indeed epinephrine. Various agencies were notified including the manufacturers of these two ampoules, and we were told that the matter would be addressed. Sadly, it appears, no real change has occurred over the past ten years. Orser and Oxorn rightfully point out that in Canada there is no federal or provincial agency that deals with these problems in drug packaging. Davies and Webb, in their accompanying editorial, remark that reporting such a problem is the first step in preventing a similar recurrence.² However, after identifying the problem, action must be taken. It appears that we are reporting the problem, we are just not acting on it. Maybe it is time that provincial and federal agencies in Canada take an interest in such labeling conflicts and similarities. Obviously the error in reading the labels on the ampoules was ours, but it is harder to put on a pair of mismatched shoes if they are a different color, shape and size!

Terrance A. Yemen MD FRCPC
Department of Anesthesia
University of Virginia Health Sciences Center
Charlottesville, Virginia 22908

REFERENCES

- 1 Orser BA, Oxorn DC. An anaesthetic drug error: minimizing the risk. *Can J Anaesth* 1994; 41: 120-4.
- 2 Davies JM, Webb RK. Adverse events in anaesthesia: the wrong drug. *Can J Anaesth* 1994; 41: 83-6.

REPLY (1)

Since reporting the case of epinephrine overdose due to a drug substitution error,¹ we have received correspondence from two other anaesthetists involved in similar cases. Dr. Yemen's comments regarding the need to do more than simply report these errors are well-founded. Recently, we met with the Labeling Committee of the Canadian Society of Hospital Pharmacists to discuss anaesthetic drug error associated with "look-alike" packaging. The development of an effective packaging system which enhances the recognition of anaesthetic drugs is not a simple task. Pharmaceutical companies are interested in measures that ensure the safe administration of their products. They require, however, convincing evidence that the need for such a system exists as well as clear guidelines as to its implementation (e.g., which drugs to include, which labeling features to modify). In order to obtain this information, a survey is being developed which will be distributed to Canadian anaesthetists.

As stated in the case report, there is no substitution for reading the label. Anaesthetists with immediate concerns about drugs with similar packaging should discuss the matter with their hospital pharmacist and consider alternative suppliers.

B.A. Orser MD FRCPC
D.C. Oxorn MD FRCPC
D.A. Yee MD FRCPC
Department of Anaesthesia
Sunnybrook Health Science Centre

REFERENCE

- 1 Orser BA, Oxorn DC. An anaesthetic drug error: minimizing the risks. *Can J Anaesth* 1994; 41: 120-4.

REPLY (2)

Yemen states that the same adverse event as detailed by Orser and Oxorn¹ occurred to a patient under his care. His comment, that "provincial and federal agencies in Canada take an interest in such labeling conflicts and similarities" is appropriate.² Some companies have made changes. For example, the packaging colours of Novolin ge Human Insulin and the beef-pork counterparts available in Canada have been changed by Connaught Novo Nordisk Inc. to be consistent with the company's international colours. (There is no world-wide standard for insulin packaging colours).

Unfortunately, in Canada there is "no federal programme for reporting drug packaging problems."¹ Absence of such a programme may reflect a low reported rate of problems or the small economic impact of such problems (i.e., both patients survived). These two factors may be the key to making changes. We would suggest that anaesthetists in Canada report not only every instance of a packaging-related drug error but also the cost of the complications suffered by the patients (i.e., ICU stay, etc.).

When they realize the incidence and magnitude of the problem the Canadian Society of Hospital Pharmacists, the Pharmaceutical Manufacturers' Association and the Canadian Drug Manufacturers' Association may understand why standard col-

our codes and labels which can be easily read are a necessity. In addition, perhaps it is time for a national body such as the Canadian Anaesthetists' Society to become involved in the problem of packaging-related drug errors in anaesthesia.

Finally, it is important to remember that drug labeling and packaging is only part of the complex patient safety issue of getting the right drug to the right patient at the right time by the right route in the right dose. There are many opportunities for human error in all stages of the action of giving a patient a drug^{3,4} and any changes made in labeling and packaging should be done in consultation with the end users.

J.M. Davies MSc MD FRPC
Department of Anaesthesia
Foothills Hospital
Calgary, Alberta
Robert K. Webb MBBS FANZCA
Australian Patient Safety Foundation
Adelaide
South Australia

REFERENCES

- 1 Orser BA, Oxorn DC. An anaesthetic drug error: minimizing the risk. *Can J Anaesth* 1994; 41: 120-4.
- 2 Yemen TA. Drug errors (Letter). *Can J Anaesth* 1994; 41: 870.
- 3 Currie M, MacKay P, Morgan C, et al. The 'wrong drug' problem in anaesthesia: an analysis of 2000 incident reports. *Anaesth Intensive Care* 1993; 21: 596-601.
- 4 Williamson JA, Webb RK, Sellen A, Runciman WB, Van der Walt JH. Human failure: an analysis of 2000 incident reports. *Anaesth Intensive Care* 1993; 21: 678-83.

Fentanyl infusions for post-thoracotomy pain management

To the Editor:

We congratulate the efforts of Dr. Baxter *et al.*¹ in their attempt to shed further light on this important and challenging topic of post-thoracotomy pain management. However, we question the validity of comparing two pairs of variables (*iv*-PCA-morphine + *iv*-fentanyl infusion vs *iv*-PCA-morphine + epidural-fentanyl infusion). It is feasible that *iv*-PCA-morphine consumption reflects an inadequately managed infusion (epidural or intravenous fentanyl) rather than demonstrating the superiority of one technique over the other. An appropriately managed infusion requires frequent patient assessment and titration. Over- or underdosing a constant infusion will add bias, and favour the more appropriately dosed infusion. As noted in Dr. Grant's editorial, fentanyl has a five-fold MEC range, therefore dosing on a per kilogram basis for a non-titrated constant infusion, is probably inappropriate. Intravenous-PCA-fentanyl vs epidural-PCA-fentanyl may be a more valid method for comparing these two techniques, rather than *iv*-PCA morphine consumption.

Larry Kahn MBChB FRCPC
Paul Jackson MD FRCPC
Department of Anaesthesia
St. Joseph's Hospital
Hamilton, Ontario L8N 1Y4

REFERENCE

- 1 Baxter AD, Laganière S, Samson B, Stewart J, Hull K, Goernert L. A comparison of lumbar epidural and intravenous fentanyl infusions for post-thoracotomy analgesia. *Can J Anaesth* 1994; 41: 184-91.

REPLY

We appreciate the interest of Drs. Kahn and Jackson in our study, and their comments about our study design.

We agree that a fixed-rate infusion regimen will not produce optimum analgesia for some patients because of variations in MEC. It is therefore important to allow the administration of additional analgesia by some means. We allowed patients to self-titrate patient-controlled morphine, as used in many other studies as an index of the efficacy of the regimen being investigated. It is possible that some patients with low MEC may have received more fentanyl than necessary with the fixed-rate infusion, and this may have produced respiratory depression. However, all patients used supplementary morphine during the study period, which we interpret as indicating inadequate analgesia at that time.

Titration of the infusion alone requires frequent assessment of the patients, may involve a latent period until changes in rate become effective, and introduces bias by involving the assessor in the "analgesic arc." "Over- or under-dosing a constant infusion" will only "add bias", as suggested by Drs. Kahn and Jackson, if the two infusion routes were treated differently (they were not in our study). Alternatively, this could have been a problem if the two routes should have been managed in different fashions and were not, but we are not aware of any evidence to suggest this.

Indeed the hypothesis to be tested by our study was that the effects of fentanyl were the same whether given by the epidural or intravenous route. We, therefore, used the same regimen by both routes: a variable dosage regimen would make interpretation of the results more difficult.

We agree that PCA administration may provide effective dose titration by both the epidural and intravenous routes. This has been used in three studies, two showing lower fentanyl requirements with epidural administration^{1,2} and the third indicating no advantage to the epidural route.³

It is noteworthy that studies using infusion titration,⁴ PCA,^{1,3} or constant infusion, as in ours, have all shown a similar pattern of poor pain control in the immediate postoperative period. Perhaps we should be looking more at other modalities, e.g., NSAIDs, intra- and/or postoperative local anaesthetics, or other opioids, rather than relying on a single agent technique for pain control in this situation.

Alan D. Baxter MB BS FRCPC FRCA
Benoit Samson MD FRCPC
Laganière Sylvie PhD
Ottawa, Ontario