intubating laryngeal mask into the trachea, the connector is detached, the intubating laryngeal mask removed, leaving the tracheal tube in place. The connector is then re-attached to the tube, and the connector can be fixed to the tube either using a super-glue or a string.

Takashi Asai MD PhD
Moriguchi City
Osaka, Japan

REFERENCES

- 1 Brain AIJ, Verghese C, Addy EV, Kapila A, Brimacombe J. The intubating laryngeal mask. II: a preliminary clinical report of a new means of intubating the trachea. *Br J Anaesth* 1997; 79: 704-9.
- 2 Brimacombe JR. Difficult airway management with the intubating laryngeal mask. *Anesth Analg* 1997; 85: 1173-5.
- 3 Asai T, Shingu K. Tracheal intubation through the intubating laryngeal mask in a patient with a fixed flexed neck and deviated larynx. *Anaesthesia* (in press).
- 4 Koga K, Asai T, Latta IP, Vaughan RS. Effect of the size of a tracheal tube and the efficacy of the use of the laryngeal mask for fibroscope-aided tracheal intubation. *Anaesthesia* 1997; 52: 131-5.
- 5 Asai T. Tracheal intubation through the laryngeal mask airway (Letter). *Anesthesiology* 1996; 85: 439.

Intubation with the LMA

To the Editor:

Joo and Rose's¹ use of the intubating LMA for difficult airways may be a safe alternative for patients who dread awake intubation, or for those who may develop acute upper airway obstruction after nerve blocks or topical anaesthesia.²

The effect of the intubating LMA on tracheal tube curvature for different 8 mm tubes at 25 and 37°C was reported by Brain *et al*³ but did not mention Joo and Rose's innovation of introducing a PCV tube with its concavity down as it enters the LMA.

Joo and Rose elected to intubate the trachea for airway management of the three difficult airways, although ventilation was achieved with the intubating LMA, the aspiration risk was low, and there was no contraindication to a conventional or flexible LMA. The use of an LMA would also have been consistent with Benumof's modification of the ASA Difficult Airway Algorithm.⁴ Only in one patient is a reason offered for intubation. Do "difficult airways" require intubation if the airway can be managed safely with an

LMA? The simple elegance of the LMA is that it allows the separation of airway priorities: effective ventilation and airway protection. If the risk of aspiration is 1: 10,000 in elective ASA I and II patients for traditional airway management or LMA,⁵ and the total lung compliance and periglottic anatomy are normal, does a difficult intubation preclude elective use of properly functioning LMA? The LMA compels us to define criteria for tracheal intubation.

Michael Beriault MD
Calgary, Alberta

REFERENCES

- 1 Joo H, Rose K. Fastrach - a new intubating laryngeal mask airway: successful use in patients with difficult airways. *Can J Anaesth* 1998; 45: 253-6.
- 2 Benumof JL. *Airway Management Principles and Practice*. Toronto: Mosby, 1996: 74-101.
- 3 Brain AIJ, Verghese C, Addy EV, Kapila A. The intubating laryngeal mask. I: Development of a new device for intubation of the trachea. *Br J Anaesth* 1997; 79: 699-703.
- 4 Benumof JL. Laryngeal mask airway and the ASA difficult airway algorithm. *Anesthesiology* 1996; 84: 686-99.
- 5 Warner MA, Warner ME, Weber JG. Clinical significance of pulmonary aspiration during the perioperative period. *Anesthesiology* 1993; 78: 56-62.

REPLY:

We believe that the angle of exit of the tracheal tube from the intubating laryngeal mask-airway (ILMA) is important not only in the success rate but possibly on post-operative sore throat. We have had high success rate (97%) in a randomized controlled study using this method.¹

Dr. Beriault's comment on the decision to intubate in the absence of aspiration risk or contraindication to LMA is cogent. The reasons for intubation in the second and third patients were subjective: the possibility of large blood loss and prolonged surgery suggested that tracheal intubation might be beneficial. However, we recognize that the LMA has been advocated as an option for patients with difficult airways if surgery is peripheral and short and when access to the airway is not compromised.² There is no evidence that the use of the LMA is any safer than tracheal intubation.

The other option in the second and third patients was to leave the ILMA as the primary airway for the surgery without tracheal intubation. However, we were concerned about possible injury to the pharyngeal mucosa after prolonged use, dental damage if the patient bit on