Correspondence

Ultrathin fiberoptic laryngoscope

To the Editor:

We read the paper by Roth *et al.*¹ with interest. The authors report successful oral tracheal intubation with the aid of a fiberoptic in 20 infants using an apnoeic relaxant technique with minimal adverse effect. We have recently completed a prospective study evaluating the 2.2 mm Olympus LF-P fibroscope (Keymed Ltd, Southend-on-Sea, Essex, UK) as an aid to oral tracheal intubation in 40 children (mean age 45) with normal airway anatomy, undergoing elective surgery using a spontaneously breathing technique with halothane.² The fibroscope was passed into the trachea in all children (mean time 69 sec) and fiberoptic intubation was successful in 30 children (mean time 113 sec). In ten children the procedure was unsuccessful; in seven the fibroscope flipped out of the trachea on attempting to railroad the tracheal tube into place, resulting in oesophageal intubation; two developed laryngospasm and in one the wrong-sized tracheal tube was initially chosen. In four children the arterial oxygen saturation decreased to <94% and in two of these children to <90%. The period of desaturation in each child lasted less than one minute.

The principle cause of failure was fibroscope displacement during railroading the tracheal tube. This was related to (1) the increased flexibility of the ultrathin fibroscope and (2) the use of preformed Portex tracheal tubes. Difficulty in railroading the tracheal tube is likely to be compounded during oro-tracheal intubation compared with naso-tracheal intubation because of the acute angulation with which the fibroscope enters the larynx. The use of a flexible armoured tracheal tube may have reduced the difficulty in railroading.^{3,4}

The conditions for fiberoptic intubation are less than ideal when using a spontaneously breathing technique with halothane. However, they are likely to be encountered during the management of the difficult paediatric airway. Therefore, we feel that this method is more useful in maintaining skills in the use of the fibroscope than a technique employing muscle-relaxants.

V.S. Sidhu FRCA
S.R. Wrigley FRCA
A.E. Black FRCA
Department of Anaesthesia

Great Ormond Street Hospital for Sick Children
Great Ormond Street
London, WC1N 3JH, UK

REFERENCES

- 1 Roth AG, Wheeler M, Stevenson GW, Hall SC. Comparison of a rigid laryngoscope with the ultrathin fiberoptic laryngoscope for tracheal intubation in infants. *Can J Anaesth* 1994; 41: 1069-73.
- 2 Wrigley SR, Black AE, Sidhu VS. A fiberoptic laryngoscope for paediatric anaesthesia. A study to evaluate the use of the 2.2 mm (LF-P) intubating fibroscope. *Anaesthesia* (in press).
- 3 Perry LB. Techniques for fiberoptic laryngoscopy in anaesthesia: guest discussion. *Anesth Analg* 1974; 53: 713-4.
- 4 Calder I. When the endotracheal tube will not pass over the flexible bronchoscope. *Anesthesiology* 1992; 77: 398.

REPLY

We thank the authors for their interest in our study and the summary of their experiences with ultrathin fiberoptic intubation. We agree that "The conditions for fiberoptic intubation are less than ideal when using a spontaneously breathing technique with halothane." In our study, we addressed the safety and efficacy of orotracheal intubation using the ultrathin fiberoptic laryngoscope in anaesthetized patients after muscle relaxation was judged to be adequate by peripheral nerve stimulation. We, too, have found using the ultrathin fiberoptic intubation technique without muscle relaxant to be a much more difficult technique, due both to the irritability of the paediatric airway (laryngospasm) and the difficulty of successfully intubating and advancing the tracheal tube into a moving glottis. In our limited experience, for ultrathin fiberoptic intubation to be successful in non-paralyzed infants, the patient must be deeply anaesthetized, with the concomitant risks of hypotension and bradycardia. For this reason, we do not routinely use that technique.

The study we reported did not address the management of children with anatomically abnormal airways but was designed to test the safety and efficacy of the ultrathin fiberoptic intubation in the child with a presumed normal airway.

Lastly, we should mention that the uncuffed, clear tracheal tubes used in our study were all made by one manufacturer (Mallinckrodt Medical, Inc., St. Louis, MO). These tubes were pre-softened in warm water to facilitate passage along the fiberoptic scope. Importantly, they were never forced into position over the fiberoptic scope, and passage was not attempted until the ultrathin fiberoptic scope was deeply positioned (near carina). With this technique, we had no difficulty in advancing the tracheal tube after intubation with the fiberoptic scope.

Andrew G. Roth MD
G.W. Stevenson MD

Steven C. Hall MD
Department of Anesthesiology
Children's Memorial Plaza
Chicago, Illinois 60614
USA

REFERENCE

- 1 Roth AG, Wheeler M, Stevenson GW, Hall SC. Comparison of a rigid laryngoscope with the ultrathin fiberoptic laryngoscope for tracheal intubation in infants. *Can J Anaesth* 1994; 41: 1069-73.

Attenuation of the pressor response to laryngoscopy – misquotation

To the Editor:

I read the article by Wig *et al.*¹ which showed an attenuating effect of calcium entry blockers upon the pressor response associated with laryngoscopy and tracheal intubation. However, they erroneously quoted our article² and stated that we could not observe any beneficial effect of verapamil. Indeed, we did find it. There is also an additional error in the statement concerning the peak effect of verapamil, which is apparent nearly one minute after *iv* injection.³⁻⁵

Toshiaki Nishikawa MD
Department of Anesthesiology
Institute of Clinical Medicine
University of Tsukuba
Tsukuba City, Ibaraki 305
Japan

REFERENCES

- 1 Wig J, Sharma M, Baichoo N, Agarwal A. Nicardipine and verapamil attenuate the pressor response to laryngoscopy and intubation. *Can J Anaesth* 1994; 41: 1185-8.
- 2 Nishikawa T, Namiki A. Attenuation of the pressor response to laryngoscopy and tracheal intubation with intravenous verapamil. *Acta Anaesthesiol Scand* 1989; 33: 232-5.
- 3 Zimpfer M, Fitzal S, Tonczar L. Verapamil as a hypotensive agent during neurolept anaesthesia. *Br J Anaesth* 1981; 53: 885-9.
- 4 Nishikawa T, Namiki A. The effects of verapamil on cerebrospinal fluid pressure in surgical patients. *Masui* 1987; 1: 132-6.
- 5 Singh BN, Roche AHG. Effects of intravenous verapamil on hemodynamics in patients with heart disease. *Am Heart J* 1977; 94: 593-9.

REPLY

Both the statements, i.e., in our article¹ and that of Nishikawa *et al.* (Ref No. 2) regarding attenuation of pressor responses are partly correct. In Nishikawa's article, tachycardia could not be attenuated. However, an increase in MAP could only be attenuated for the first 5-10 min. Later, the difference in the rise in MAP in both the control and verapamil-treated groups was insignificant. We did not fully elucidate this fact. Also, we apologize for the second error and will be more careful in future.

Jyotsna Wig MD
Chandigarh
India

Laryngoscopy and arrhythmias in children

To the Editor:

We would like to report our observations involving the use of halothane during induction, laryngoscopy, and tracheal intubation in children with supraventricular tachycardia (SVT) syndromes. With the establishment of a paediatric cardiac electrophysiological study (EPS) diagnostic laboratory at our institution, that included radiofrequency ablation (RFA) procedures when appropriate, the anaesthesia department was asked to provide general anaesthesia for these children. This was due to the anticipated prolonged nature of the cases (4-12 hr), the potential for complications (e.g., SVT requiring cardioversion) and the desire to have a motionless, stable child during placement of the multiple catheters.

Since it was necessary to enable the cardiologist to elicit SVT for both diagnostic and therapeutic reasons, we initially selected halothane as our primary anaesthetic due to its reputation for arrhythmogenicity as well as our familiarity with the agent in routine paediatric anaesthetic practice. Dealing with a patient group that sought RFA due to life-threatening arrhythmias, medication failures or complications, or desire for a definitive correction, we were concerned about eliciting SVT during induction and intubation. We selected a protocol consisting of benzodiazepine premedication (diazepam 0.2 mg · kg⁻¹), placement of an *iv* catheter upon arrival in the catheterization laboratory, and *iv* induction with thiopentone 4 mg · kg⁻¹, vecuronium 0.1 mg · kg⁻¹, and then inhalational halothane, 1.5% end-tidal concentration, N₂O 3 L · min⁻¹ and O₂ 2 L · min⁻¹ with PETCO₂ 35-40 mmHg. Continuous monitoring for SVT was recorded, with hardcopy printouts prior to induction, after induction with achievement of end-tidal halothane concentrations of 1.5% for three minutes, and again immediately after laryngoscopy and intubation.

No arrhythmias were observed during induction, lar-