lesions (nerve root demyelination and Wallerian degeneration) were observed in all groups.³ However, others have studied the neurotoxicity of intrathecal midazolam.^{4,5} We first observed a high incidence of lesions in rabbits,⁴ and our findings were confirmed by Svensson *et al.*, who characterised the nature of the toxicity.⁵ It is not possible to promote intrathecal midazolam in man since neurotoxicity has been reported in rabbits⁴ and in rats.⁵ Even if neurotoxicity of epidurally administered drug is attenuated, we must keep in mind that the risk of unintended intrathecal injection of epidurally administered agent is not negligible.

Jean-Marc Malinovsky MD
Nantes, France

REFERENCES

- 1 Bahar M, Cohen ML, Grinshpon Y, Chanimov M. Spinal anaesthesia with midazolam in the rat. *Can J Anaesth* 1997; 44: 208–15.
- 2 Goodchild CS, Noble J. The effects of intrathecal midazolam on sympathetic nervous system reflexes in man – a pilot study. *Br J Clin Pharmacol* 1987; 23: 279–85.
- 3 Serrao JM, Mackenzie JM, Goodchild CS, Gent JP. Intrathecal midazolam in the rat: an investigation of possible neurotoxic effects. *Eur J Anaesthesiol* 1990; 7: 115–22.
- 4 Malinovsky J-M, Cozian A, Lepage J-Y, Mussini J-M, Pinaud M, Souron R. Ketamine and midazolam neurotoxicity in the rabbit. *Anesthesiology* 1991; 75: 91–7.
- 5 Svensson BA, Welin M, Gordh T Jr, Westman J. Chronic subarachnoid midazolam (Dormicum) in the rat. Morphologic evidence of spinal cord neurotoxicity. *Reg Anesth* 1995; 20: 426–34.

REPLY

We are pleased to respond to Malinovsky's criticism of our study.¹ We have just completed a neurotoxicity study on intrathecal midazolam in the same animal model which awaits acceptance for publication. We examined whether midazolam injected intrathecally, alone or combined with fentanyl, on 15 occasions over one month produced neurotoxic injury as judged by the animal's neurological behaviour and by histological examination, by light microscopy, of the excised spinal cord and paraspinal tissues on sacrifice after five weeks. The white matter showed a spongiose appearance, but no evidence of demyelination. The neurones of both the anterior and posterior horns of grey matter showed cytoplasmic vacuolation of varying degree, but no differences in these changes were observed among groups that received midazolam alone, midazolam plus fentanyl, and those given intrathecal lidocaine alone or fentanyl alone. After each intrathecal injection

all animals recovered fully and remained awake, mobile and ate and drank normally. They continued to do so until the end of the study five weeks later.

The sophisticated neurotoxicity studies by Malinovsky *et al.*² and by Svensson *et al.*³ employing morphometric and ultrastructural endpoints after chronic subarachnoid midazolam injection, showed subtle changes in the midazolam group of animals, compared with those who received saline or lidocaine. None of the animals presented obvious neurological impairment or behavioural disturbances.

We fully endorse Yaksh's admonition that it is necessary to rule out potential neurotoxic effects of novel drug combinations in laboratory animals before administering such combinations to patients.⁴

M. Bahar MD,
M. Cohen MB CHB,
Y. Grinshpon MD,
M. Chanimov MD
Zerifin, Israel

REFERENCES

- 1 Bahar M, Cohen ML, Grinshpon Y, Chanimov M. Spinal anaesthesia with midazolam in the rat. *Can J Anaesth* 1997; 44: 208–15.
- 2 Malinovsky J-M, Cozian A, Lepage J-Y, Mussini J-M, Pinaud M, Souron R. Ketamine and midazolam neurotoxicity in the rabbit. *Anesthesiology* 1991; 75: 91–7.
- 3 Svensson BA, Welin M, Gordh T Jr, Westman J. Chronic subarachnoid midazolam (Dormicum) in the rat. Morphologic evidence of spinal cord neurotoxicity. *Reg Anesth* 1995; 20: 426–34.
- 4 Yaksh TL, Collins JG. Studies in animals should precede human use of spinally administered drugs. *Anesthesiology* 1989; 70: 4–6.

Protamine and its cardiovascular effects

To the Editor:

Backman, Gilron and Robbins' case report makes for interesting observation.¹ They rightly discount the mechanism of acute cardiovascular events following protamine injection as due to a vaso-vagal reaction via reinnervation but, rather, suggest the possible causes as reflex decrease in sympathetic output secondary to stimulation of mechano-receptors, reduced sino-atrial node perfusion, or possible cardiac ischaemia.

Although the cause of the well reported cardiovascular events are uncertain, animal models have indicated possible mechanisms of action. Protamine has been shown to interfere with mitochondrial respiration *in vitro*² and, in dogs, reduces *in vivo* oxygen