ing comment: "Certainly it is apparent that every child deserves special consideration, and there is little place in paediatric anaesthesia practice for the use of routine schedules of pre-operative drugs."⁶

Victor Faria Blanc MD FRCP(C)
Hôpital Ste-Justine and
Université de Montréal
Montreal

REFERENCES

- 1 Steward DJ. Anticholinergic premedication for infants and children. Editorial. *Can Anaesth Soc J* 1983; 30: 325-6.
- 2 Blanc VF, Hardy JF, Milot J, Jacob JL. The oculo-cardiac reflex: a graphic and statistical analysis in infants and children. *Can Anaesth Soc J* 1983; 30: 360-9.
- 3 Donlon JV Jr. Anesthesia for eye, ear, nose and throat surgery. In Miller RD, Editor. *Anesthesia* Vol. 2, Chpt. 39: 1273, Churchill Livingstone, N.Y., 1981.
- 4 Smith RM. Anesthesia for infants and children - 4th edit. C.V. Mosby, St. Louis, 1980.
- 5 Steward DJ. Manual of pediatric anesthesia. Longman Canada Ltd., Don Mills, Ontario. 1979.
- 6 Steward DJ. Psychological aspects of anaesthetising infants and children. In Steward DJ, Editor. *Some aspects of paediatric anaesthesia*. Excerpta Medica, Amsterdam 1982.

REPLY

I am sorry that Dr. Blanc is upset because, in my editorial, I failed to stress what he feels is important, i.e., the need for a better understanding of the oculo-cardiac reflex (OCR). Indeed, I was more concerned with examining the use of anticholinergics for paediatric patients.

From their study, Blanc et al. concluded "intramuscular atropine associated with gentle (or more physiological) manipulation of the extra-ocular muscles can reduce the intensity of the OCR, as well as the occurrence of cardiac arrhythmias, to very acceptable levels and can provide a way of avoiding the potential risks inherent to the routine use of intravenous anticholinergics." I do not believe that this conclusion is justified by a study of 49 infants and children. The vast majority of investigators have concluded that intravenous atropine is the surest and safest method to block the OCR. The "risks" of intravenous anticholinergics in paediatric patients are not at all proven. There are no documented cases of serious complications following IV anticholinergics in healthy children.

Even the suggestion that controlled ventilation may modify the incidence of the OCR is at best controversial. A very recent report suggests that, during halothane anaesthesia, the use of controlled ventilation did not reduce the occurrence of the OCR.¹

My advice must remain the same: to avoid risk of serious complication due to the OCR, administer an intravenous anticholinergic before manipulation of the extra-ocular muscles commences.

Dr. Blanc and his colleagues have contributed significantly to our understanding of the mechanisms of the OCR. Their results, however, do not of necessity dictate that we should change the clinical management of our patients.

D.J. Steward MB FRCP(C)
Anaesthetist-in-Chief
The Hospital for Sick Children
Toronto

REFERENCE

- 1 Forestner JE, Imbrecht P. Controlled respiration does not inhibit oculocardiac reflex during strabismus surgery in children. *Anesthesiol* 1983; 59: A457.