TABLE Suggested length of time before anesthetizing a patient following coronary stent replacement

Number of weeks	n	(%)
0 weeks	13	3.8
1 week	12	3.5
2 weeks	87	25.3
3 weeks	13	3.8
4 weeks	127	36.9
5 weeks	52	15.1
6 weeks	13	3.8
7 weeks	2	0.6
8 weeks	7	2.0
12 weeks	16	4.7
24 weeks	2	0.6
Total	344	

often faced with a dilemma about how long to wait between stent insertion and a subsequent surgical procedure. Patients may be placed at risk by either a too short or too long waiting period. The American Heart Association along with the American College of Cardiologists recommend a minimum wait of two weeks and suggest an ideal waiting period of four to six weeks, although these guidelines are not based on randomized controlled trials.¹ It is unclear whether Canadian anesthesiologists are aware of these guidelines. The aim of this survey was to describe the views of practicing anesthesiologists on the waiting times for elective surgery following insertion of a coronary stent.

A postal survey (along with a reminder eight weeks later) was sent to 2,167 Canadian anesthesiologists in March, 2003 seeking their views about the appropriate waiting time for unrelated elective surgery following insertion of a coronary stent. Responders were presented with a clinical scenario (a 59-yr-old male with a prior stent insertion, no comorbidities, who required an urgent aneurysm repair) and asked to report a recommended waiting time for three different locations (left anterior descending, circumflex or right coronary arteries). They were also asked about the existence of a policy on this issue in their departments.

The response rate was 46%. Most departments of anesthesiology (97%) do not have an explicit policy regarding the appropriate waiting time between coronary stent insertion and elective surgery. Half of the responders believed their departments ought to have such a policy. While there was no consensus about what this policy ought to be, the majority of those who wanted a policy felt that the minimum waiting time should be four weeks following surgery (Table). Few anesthesiologists felt that the minimum waiting time should be less than four weeks. A substantial pro-

portion (63%) of Canadian anesthesiologists were unaware of published or unpublished literature, including published U.S. guidelines, about the appropriate length of time between the insertion of a stent and a subsequent surgical procedure.

The implications of this study are: 1) Canadian departments of anesthesiology consider implementing an explicit policy that sets the waiting time for unrelated, elective surgery following coronary stent insertion, to four weeks or greater; and 2) anesthesiologists review published U.S. recommendations.¹

Lindsey Patterson FRCA

Duncan Hunter PhD

Andrea Mann BSc

Queen's University, Kingston, Canada

E-mail: ljp@post.queensu.ca

Source of support: Queen's University Faculty of Health Sciences Research Initiation Grant.

Acknowledgements

Drs. Ted Ashbury, David Mark, Brian Milne, Joel Parlow, and Chris Simpson for reviewing the questionnaire. The Royal College of Physicians and Surgeons of Canada and the Canadian Anesthesiologists' Society for providing the mailing list. The anesthesiologists who took the time to complete the questionnaire. The Queen's University Faculty of Health Sciences Research Initiation Grant for funding this research.

Reference

- 1 Eagle KA, Berger PB, Calkins H, et al. ACC/AHA Guideline Update for Perioperative Cardiovascular Evaluation for Noncardiac Surgery—Executive Summary. A report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Committee to update the 1996 Guidelines on Perioperative Cardiovascular Evaluation for Noncardiac Surgery). *Anesth Analg* 2002; 94: 1052–64.

S100 β and postcardiac surgery neurological dysfunction: reasons to disregard any link

To the Editor:

Dr. Hall's editorial¹ in reference to the work by Kanbak *et al.*,² eloquently outlines the pitfalls of using S100 β as a surrogate marker of postcardiac surgery neurologic dysfunction. He explains why S100 β , a

glial protein long considered to be a specific marker of brain injury, may be elevated in the setting of cardiac surgery when no apparent neurologic injury (as described by Kanbak *et al.*)² has occurred. The reason for this relationship (i.e., lack thereof) relates to apparent sources for S100 β . In addition to glial cells, it has been found in the mediastinal blood and when retransfused to the patient via the cardiectomy suction, subsequently leads to significant elevations in serum S100 β . However, to further clarify this point, one needs to elaborate why an erroneous relationship between S100 β and neurologic injury has repeatedly been made in the literature. It is commonly accepted that some of the particulate lipid-containing debris and blood aspirated by the cardiectomy suction from the mediastinum and pericardium during surgery makes its way back into the venous reservoir and passes through the cardiopulmonary bypass apparatus thus serving as a source of cerebral microemboli.³ In this regard, if the mediastinal drainage can cause injury by increasing cerebral embolization, and in an unrelated epiphenomenal manner also increases serum S100 β due to its apparent high S100 β content, an erroneous S100 β /cerebral injury relationship is realized.⁴ Importantly, however, further light has recently been shed on the source of this mediastinal S100 β . Fazio *et al.*, have now demonstrated that there is in fact no extracerebral S100 β source in the mediastinum and that the error lies in the lack of S100 β specificity of conventional assays.⁵ These investigators have elegantly demonstrated that it is other large molecular weight protein contaminants (such as haptoglobin I precursor) that are falsely detected by conventional assays that are responsible for these elevated, although erroneous, S100 β levels. These recent publications reinforce how, at least in the setting of cardiac surgery, S100 β is clearly not ready yet for prime time as Dr. Hall states, and furthermore, likely never will be.⁶

Hilary P. Grocott MD FRCPC
Duke University Medical Center, Durham, USA
E-mail: h.grocott@duke.edu

References

- 1 Hall RI. Serum S-100 β protein and postoperative neurological dysfunction--ready for prime time? *Can J Anesth* 2004; 51: 645-8.
- 2 Kanbak M, Saricaoglu F, Avci A, Ocal T, Koray Z, Aypar U. Propofol offers no advantage over isoflurane anesthesia for cerebral protection during cardiopulmonary bypass: a preliminary study of S-100 β protein levels. *Can J Anesth* 2004; 51: 712-7.
- 3 Brooker RF, Brown WR, Moody DM, *et al.* Cardiectomy suction: a major source of brain lipid emboli during cardiopulmonary bypass. *Ann Thorac Surg* 1998; 65: 1651-5.
- 4 Grocott HP, Arrowsmith JE. Serum S100 protein as a marker of cerebral damage during cardiac surgery. *Br J Anaesth* 2001; 86: 289-90.
- 5 Fazio V, Bhudia SK, Marchi N, Aumayr B, Janigro D. Peripheral detection of S100 β during cardiothoracic surgery: what are we really measuring? *Ann Thorac Surg* 2004; 78: 46-52; discussion 52-3.
- 6 Grocott HP. Invited commentary. *Ann Thorac Surg* 2004; 78: 52-3.

REPLY:

We welcome the opportunity to respond to the letter of Grocott. The points raised regarding extra cranial sources of S100 β protein are valid. As the biological half-life of S100 β protein is less than one hour.¹ We selected a sampling time of 24 hr after cardiopulmonary bypass (CPB). Georgiadis and coworkers found that S100 β protein levels at 24 hr after CPB have a sensitivity and specificity of approximately 90% and 97% respectively.² As Shaaban Ali mentioned, unless there is a ongoing auto transfusion from chest tubes, secondary or late release pattern of S100 β protein (15-48 hr) in the post-operative period is a sign of further neurological injury.³

Fazio *et al.*⁴ have demonstrated the limitations associated with the use of S100 β protein for the evaluation of brain damage and patients' outcome during surgical procedures but they concluded that currently available methods used to measure serum S100 β protein are still valid for the evaluation of neurological disorders associated with brain damage. Filtering or immunoprecipitation of samples may be required to completely rule out cross contamination.

In addition to these, our study aimed to investigate the effect of anesthetics on the S100 β protein, and our groups were standardized for the surgical technique and measurements.

Prospective, large-scale, multicentre studies will be necessary to determine the exact origins of S100 β protein.

Meral Kanbak MD

Fatma Saricaoglu MD

Turgay Ocal MD

Hacettepe University, Ankara, Turkey

E-mail: orhankan@ttnet.net.tr

References

- 1 Jonsson H, Jonsson P, Alling C, Backstrom M, Bergh C, Blomquist S. S100 β after coronary artery surgery: release pattern, source of contamination, and relation to neu-

ropsychological outcome. *Ann Thorac Surg* 1999; 68: 2202–8.

- 2 Georgiadis D, Berger A, Kowatschev E, et al. Predictive value of S-100 β and neuron-specific enolase serum levels for adverse neurologic outcome after cardiac surgery. *J Thorac Cardiovasc Surg* 2000; 119: 138–47.
- 3 Shaaban Ali M, Harmer M, Vaughan RS. Serum S100 protein as a marker of cerebral damage during cardiac surgery (Letter, reply). *Br J Anaesth* 2001; 86: 289–90.
- 4 Fazio V, Bhudia SK, Marchi N, Aumayr B, Janigro D. Peripheral detection of S100 β during cardiothoracic surgery: what are we really measuring? *Ann Thorac Surg* 2004; 78: 46–53.

Intrathoracic positioning of a thoracic epidural catheter inserted via the median approach

To the Editor:

Thoracic epidural catheterization for perioperative anesthesia is well-established and used frequently in combination with general anesthesia for abdominal and thoracoabdominal surgery.¹ A possible complication related to technique, the accidental puncture of the pleural cavity or intrathoracic migration of the catheter has been reported rarely, albeit by the paramedian approach.² We report a case of intrathoracic placement of a thoracic epidural catheter inserted via the median approach.

A thoracic epidural catheter was inserted and advanced 6.5 cm into the epidural space using a mid-line approach at the T8–9 level in a 71-yr-old woman undergoing elective esophagectomy and high thoracic esophagogastronomy. The epidural space was identified easily on the first attempt at 7 cm by the loss-of-resistance technique. No pain, shortness of breath, cough or shoulder pain and no neurological signs were observed during puncture. During surgery, the catheter was seen to lie in the right pleural cavity. The position of the catheter was documented by a photograph and the catheter was withdrawn immediately. As the photograph demonstrates, the catheter penetrated the parietal pleura close to the spine. The 5.5-cm mark on the catheter cannot be seen in the pleural cavity, which suggests that the point of loss-of-resistance (and therefore the tip of the Tuohy needle) was at least 1 cm from the parietal pleura. This indicates that the pleural penetration was likely made by the catheter and not by the Tuohy needle.

FIGURE View into the right pleural cavity. The catheter penetrates the pleura lateral to the spinal column. The tip of the catheter (arrow) is visible while the 5.5-cm mark is not.

Accidental pleural puncture or the intrathoracic placement of catheters via a thoracic epidural are a rare, but potentially dangerous complication, especially in procedures with thoracotomy and one-lung-ventilation on the contralateral side. The median approach is thought to be less likely to produce pneumothorax and all reported cases to date occurred using a paramedian approach. This is the first report of pleural puncture by a thoracic epidural catheter using a median approach.^{3,4} Further investigation is necessary to demonstrate if pleural puncture in general is more likely to occur than previously thought and is more likely when using the paramedian approach.

One must assume that poor or ineffective analgesia, so-called "failed" epidurals may be due to misplacement and this includes interpleural position.^{2,5} The correct insertion of catheters into the epidural space can thereby only be verified by demonstrating a neural block and rare complications should be kept in mind when performing a thoracic epidural anesthesia.⁵

Britta Patermann MD

John Lynch FFARCS

Paul Schneider MD

Christoph Weigand MD

Sandra Kampe MD

University of Cologne, Cologne, Germany

E-mail: Sandra.Kampe@medizin.uni-koeln.de

References

- 1 Moraca RJ, Sheldon DG, Thirlby RC. The role of epidural anesthesia and analgesia in surgical practice.