outcome of the infraclavicular brachial plexus blocks. At our institution, instead of making use of “three factors,” we utilize one common endpoint to ensure the success of the blocks. It is the deposition of the local anesthetic (LA) around the brachial plexus cords. We believe that having the LA “bathe” the nerves is important in predicting the success of the blocks. The U-shaped distribution of LA and the anterior displacement of the artery are often seen as a result of the correct deposition of the LA. Fascial click, however, is sometimes felt twice during the blocks. One click is always felt as the needle pierces the neurovascular sheath, and the second click is sometimes felt as it passes through the septum.

Dr. Lévesque writes, “... we need to think in terms of the diffusion compartment, instead of focusing on pure needle-to-nerve distance”. We, too, feel that observing the appropriate LA spread is far more important than having the block needle near the nerve at the beginning of the block. Sinha et al.¹ have previously reported that the presence of nerve stimulation had no impact on the success of the resulting blocks. Not only does the ultrasound technology allow us to locate the target structures, it also allows us to direct the needle and to observe the real-time LA spread. Ultimately, this know-how should lead to safer and more effective blocks.

Maki Morimoto MD

Jovan Popovic MD

Jung T. Kim MD

Harald Kiamzon MD

Andrew D. Rosenberg MD

NYU Medical Center, New York, USA

E-mail: maki.morimoto@med.nyu.edu

Reference

- 1 Sinha SK, Abrams JH, Weller RS. Ultrasound-guided interscalene needle placement produces successful anesthesia regardless of motor stimulation above or below 0.5 mA. *Anesth Analg* 2007; 105: 848–52.

Hypertrophic cardiomyopathy – a matter of genes

To the Editor:

We write to comment on the *Images in Anesthesia* feature by Fayad,¹ which describes a patient with undiagnosed hypertrophic obstructive cardiomyopathy, who developed hypotension during anesthesia. Because the patient had a long-standing history of hypertension, we suggest that the cardiac condition in this

FIGURE A Diastolic image of a steady state free precession (SSFP) sequence in the four-chamber view of the above-mentioned patient. The septal hypertrophy can be clearly seen (asterisk).

patient may not have been secondary to hypertrophic (obstructive) cardiomyopathy.

Hypertrophic cardiomyopathy (HCM) is a genetic disorder. Diagnosis usually requires the exclusion of other causes of left ventricular hypertrophy — in particular, arterial hypertension, and aortic stenosis.² To date, some 400 mutations, in at least 11 genes, have been shown to cause HCM, with an autosomal dominant pattern of inheritance. The phenotype of HCM is highly variable, ranging from asymptomatic cases, to patients who are severely limited by dyspnea and angina. Cardiac hypertrophy in affected individuals can be restricted to the interventricular septum, or to virtually any part of the ventricle. The disease can also present as concentric hypertrophy. Accordingly, left ventricular outflow tract (LVOT) obstruction is not a mandatory feature of HCM. Many patients with HCM do not have LVOT obstruction at rest, but develop LVOT gradients during exercise (latent hypertrophy). Alternatively, dynamic LVOT obstruction may be restricted in patients with HCM to conditions involving a hyperdynamic circulation (e.g., secondary to anemia). Dynamic LVOT obstruction can also be precipitated by drugs that reduce cardiac afterload, thus increasing the gradient between the left ventricle and the aorta. However, a diagnosis of LVOT obstruction is not pathognomonic for HCM. Patients with long-standing and/or pronounced arterial hypertension may also present with LVOT obstruction. Since LVOT obstruction is a dynamic phenomenon, even patients with structurally normal hearts may develop LVOT obstruction.³

FIGURE B Corresponding diastolic image in the three-chamber view (3CV): the ascending aorta (AO), left ventricular cavity (LV) and left atrium (LA) can be identified.

FIGURE C Mid-systolic image in the three-chamber view (3CV): turbulent flow in the left ventricular outflow tract (arrowhead) and systolic anterior movement of the anterior leaflet of the mitral valve (SAM-phenomenon) due to septal hypertrophy (asterisk). The dynamic LVOT gradient and mitral regurgitation are best seen on a cine-loop of the 3CV (available as Additional Material at www.cja-jca.org).

Although the clinical presentation of HCM and hypertensive heart disease may be similar, the clinical and therapeutic consequences differ. Whereas angiotensin converting enzyme inhibitors, angiotensin receptor antagonists, or diuretics are effective drugs to regulate high blood pressure, they are not indicated for most patients with HCM. Furthermore,

HCM patients should be discouraged from participating in competitive sports. They need regular and careful evaluation for specific risk factors of sudden cardiac death.⁴ Since sudden cardiac death may occur in asymptomatic HCM mutation carriers, regular clinical screening of close relatives is strongly advised. None of these measures is usually required in patients with arterial hypertension.

To illustrate the existence of LVOT obstruction in the absence of HCM, we describe the case of a 37-yr-old patient with a nine-year history of renal failure due to IgA nephropathy, associated with marked arterial hypertension. The patient had received a kidney transplant seven years previously, but transplant failure developed three years later. Successful kidney re-transplantation was done only recently. One year before kidney re-transplantation, the patient developed severe exertional dyspnea and angina. Marked asymmetrical left ventricular hypertrophy and resting LVOT obstruction resembling HCM were observed by echocardiography, and the cardiac magnetic resonance imaging (Figure). Cardiac examination of her first degree relatives excluded the presence/coexistence of familial HCM. As the patient's symptoms could not be controlled by medical therapy, she underwent transcatheter ablation of a septal hypertrophy, which resulted in dramatic improvement of her symptoms.

This case demonstrates that hypertensive left ventricular hypertrophy can present with typical features of HCM. We suggest that one should consider the possible coexistence of HCM and arterial hypertension, only in those patients harbouring a causative mutation in one of the known HCM disease genes, or in patients with a family history of HCM.

Christian Geier MD

Thomas Elgeti MD

Cemil Ozcelik MD

Charité University Hospital Berlin, Berlin, Germany

E-mail: christian.geier@charite.de

Conflicts of interests and sources of support: None declared.

Accepted for publication February 20, 2008.

References

- 1 Fayad A. Left ventricular outflow obstruction in a patient with undiagnosed hypertrophic obstructive cardiomyopathy. *Can J Anesth* 2007; 54: 1019–20.
- 2 Maron BJ, Towbin JA, Thiene G, et al. Contemporary definitions and classification of the cardiomyopathies: an American Heart Association Scientific Statement from the Council on Clinical Cardiology, Heart Failure

and Transplantation Committee; Quality of Care and Outcomes Research and Functional Genomics and Translational Biology Interdisciplinary Working Groups; and Council on Epidemiology and Prevention. *Circulation* 2006; 113: 1807–16.

- 3 Mingo S, Benedicto A, Jimenez MC, Perez MA, Montero M. Dynamic left ventricular outflow tract obstruction secondary to catecholamine excess in a normal ventricle. *Int J Cardiol* 2006; 112: 393–6.
- 4 Maron BJ, McKenna WJ, Danielson GK, et al. American College of Cardiology/European Society of Cardiology clinical expert consensus document on hypertrophic cardiomyopathy. A report of the American College of Cardiology Foundation Task Force on Clinical Expert Consensus Documents and the European Society of Cardiology Committee for Practice Guidelines. *J Am Coll Cardiol* 2003; 42: 1687–713.

Reply:

I thank Drs. Geier et al. for their interest in the recently published *Images in Anesthesia* feature.¹ Dr. Geier is correct in detailing the etiology and pathophysiology of hypertrophic cardiomyopathy (HCM). The case report however, does not in any way imply that the cause for the left ventricular outflow tract (LVOT) obstruction was the presence of the long standing hypertension. It rather highlighted the positive findings of the clinical history and physical exam, with the main emphasis being on the echocardiography images. As mentioned, most patients with HCM are asymptomatic.² However, dynamic LVOT obstruction can occur in patients with normal ventricles,³ as well as in patients with hypertensive heart disease. In context of an *Images in Anesthesia* feature, the main purpose was to demonstrate the echocardiographic images of LVOT obstruction, and to highlight the importance of the intraoperative echocardiography as a monitor in confirming the diagnosis.

Ashraf Fayad MD FRCPC
University of Ottawa, Ottawa, Canada
E-mail: afayad@ottawahospital.on.ca

References

- 1 Fayad A. Left ventricular outflow obstruction in a patient with undiagnosed hypertrophic cardiomyopathy. *Can J Anesth* 2007; 54: 1019–20.
- 2 Spirito P, Seidman CE, McKenna WJ, Maron BJ. The management of hypertrophic cardiomyopathy. *N Engl J Med* 1997; 336: 775–85.
- 3 Mingo S, Benedicto A, Jimenez MC, Perez MA, Montero

M. Dynamic left ventricular outflow tract obstruction secondary to catecholamine excess in a normal ventricle. *Int J Cardiol* 2006; 112: 393–6.

Methylene blue is a potent monoamine oxidase inhibitor

To the Editor:

I was pleased to read Dr. Ng's case report, for which he sought my expertise and assistance, but sorry he did not avail himself of our results, that have established the monoamine oxidase inhibitors (MAOI) potency of methylene blue, published in September 2007.¹

It is therefore worthwhile adding the 'final chapter' to this interesting story for your readers, and rectifying some particulars that are unclear or inexact in his discussion. My initial report in 2006² postulated that the severity of the serotonin toxicity symptoms in these cases was such as to indicate methylene blue *must* be acting as an MAOI. I stated '... further corroboration and quantification of methylene blue's potency (as an MAOI) is in progress.' That paper has confirmed the prediction by demonstrating experimentally, that it is a potent tight binding MAO-A inhibitor at nanomolar concentrations: i.e., it is ~ 100 times more potent than moclobemide.¹ There can now be no doubt about the cause effect relationship of drugs [methylene blue + selective serotonin reuptake inhibitors (SSRIs)] to symptoms (serotonin toxicity).

Furthermore, serotonin toxicity is predictable because the spectrum concept explains it as a continuum of increasing effects, not an idiosyncratic reaction like neuroleptic malignant syndrome (this is why it is clearer to stick to the term serotonin toxicity, and avoid serotonin syndrome). This, and other aspects of serotonin toxicity, are covered in reviews.^{3,4} Different classes of drugs exhibit distinct degrees to which they can elevate serotonin.⁵ Thus, over-doses of SSRIs produce only mild to moderate degrees of serotonin toxicity, whereas combinations of monoamine oxidase inhibitors, (including reversible inhibitors of monoamine oxidase-A like moclobemide) with any serotonin reuptake inhibitor, *predictably and frequently* precipitate severe serotonergic side effects that can progress to frank serotonin toxicity (depending on dose).

I do endorse Dr. Ng's point 'This illustrates the importance of a high level of clinical suspicion and an appropriately focussed assessment.' But it is also crucial to link that idea to the spectrum concept, and to appreciate that increasing serotonergic side effects and serotonin toxicity is likely, not just possible, with