Short-acting neuromuscular blocking drugs allow better control of OR time

To the Editor:

I read with interest the article by Dahaba and colleagues comparing rocuronium and mivacurium equi-lasting doses in patients undergoing laparoscopic gynecological surgery.¹ As the authors stated, the duration of this type of surgery is extremely variable. Since mivacurium is hydrolyzed by plasma cholinesterase the recovery time from the blockade does not depend on the total cumulative dose by either repeated doses or continuous infusion.² It only depends on the presence of normal functioning plasma cholinesterase. This feature makes it unique especially when the duration of surgery is unpredictable. Stopping the infusion or maintenance dose administration at any time will be followed immediately by a fast recovery. In contrast, if the surgery is concluded immediately after a maintenance dose of rocuronium, a longer time will be needed for full recovery even if a reversal drug is given. Reid *et al.* found that after neostigmine reversal of rocuronium-induced neuromuscular blockade when T1 was 25%, only 14 out of 40 patients attained a train-of-four ratio of 0.8 within 15 min of reversal administration.³ The difference in drug acquisition costs according to the authors was only 2 /patient in favour of rocuronium but the difference in recovery times (six minutes) was 45 /patient in favour of mivacurium. Even if we agree on the difference the authors reported in the recovery times, mivacurium use still costs 43 less/patient. If we add to this the frustration, that may result from delaying the operating room (OR) schedule, which will affect the next patient in the schedule, the surgeon, the nursing team as well as the anesthesiologist himself, we can conclude that mivacurium allows better control of OR time avoiding delays in patient recovery and its associated economic and personnel implications.

Mohammad El-Orbany MD
Chicago, Illinois

References

- 1 Dahaba AA, Schweitzer E, Fitzgerald RD, Schwarz S. Equi-lasting doses of rocuronium, compared to mivacurium, result in improved neuromuscular blockade in patients undergoing gynecological laparoscopy. *Can J Anesth* 2001; 48: 1084–90.
- 2 Savarese JJ, Lien CA, Belmont MR, Rubin L. The clinical and basic pharmacology of mivacurium: a short-acting nondepolarizing benzylisoquinolinium diester neuromuscular blocking drug. *Acta Anaesthesiol Scand*

Suppl 1995; 106: 18–22.

- 3 Reid JE, Breslin DS, Mirakhor RK, Hayes AH. Neostigmine antagonism of rocuronium block during anesthesia with sevoflurane, isoflurane or propofol. *Can J Anesth* 2001; 48: 351–5.

REPLY:

One out of 40 Caucasians are heterozygous for the atypical and the silent plasma cholinesterase gene, while one out of 2,800 are homozygous for the atypical gene. In these patients, neuromuscular blockade with mivacurium will be prolonged markedly, may last up to ten hours¹ and could prove hard to reverse even with several doses of neostigmine.² This would represent, in our hospital, an additional postanesthesia care unit or intensive care unit cost of \$4,500–\$9,000. It is likely that most anesthesiologists, in the course of their career, will encounter several cases of prolonged response to mivacurium (5–7% of patients of an unselected population).³ Needless to say that accelerating the reversal of such a block using plasma cholinesterase derived from pooled human blood, with the inherent risk of infectious disease transmission, is not desirable. Screening all patients for abnormal plasma cholinesterase is not a practical solution as it is too expensive and time-consuming.⁴

In our discussion, we presented drug acquisition costs separately from the cost of running our operating room, which would obviously vary greatly between different hospitals. We feel strongly that decisions regarding time and cost should be left entirely to the discretion of each anesthesiologist who might, for instance, choose to avoid the costs associated with delayed mivacurium recovery. Others would choose to resort to a neuromuscular blocking drug infusion, as Dr. El-Orbany mentioned, or reversal of a residual mivacurium block although it might increase the risk of postoperative nausea and vomiting in gynecological laparoscopy.⁵ However, these considerations were beyond the scope and design of our study.

Ashraf A. Dahaba MD MSc
Graz, Austria

References

- 1 Viby-Mogensen J. Mivacurium block in patients with abnormal plasma cholinesterase. *Acta Anaesthesiol Scand* 1995; 39: 1001–2.
- 2 Petersen RS, Bailey PL, Kalameghian R, Ashwood ER. Prolonged neuromuscular block after mivacurium. *Anesth Analg* 1993; 76: 194–6.
- 3 Hanel HK, Viby-Mogensen J, de Muckadeel OB. Serum cholinesterase variants in the Danish population. *Acta Anaesthesiol Scand* 1978; 22: 505–7.
- 4 Maddineni VR, Mirakhor RK. Prolonged neuromus-

cular block following mivacurium. *Anesthesiology* 1993; 78: 1181-4.

- 5 Lovstad RZ, Thagaard KS, Berner NS, Raeder JC. Neostigmine 50 microg kg(-1) with glycopyrrolate increases postoperative nausea in women after laparoscopic gynaecological surgery. *Acta Anaesthesiol Scand* 2001; 45: 495-500.

The incidence of failed spinal anesthesia, postdural puncture headache and backache is similar with Atraucan and Whitacre spinal needles

To the Editor:

The Atraucan (AT) dual-bevelled spinal needle, which has been shown, *in vitro*, to cause less cerebral spinal fluid (CSF) leakage than comparable Quincke, Sprotte and Whitacre (WH) spinal needles,^{1,2} has been available in the United States since 1993. Three clinical studies³⁻⁵ compared the success or complication rate of AT spinal needles to pencil-point spinal needles. Vallejo *et al.*⁵ reported a higher incidence of epidural blood patch and postdural puncture headache (PDPH) associated with 26-gauge AT spinal needles compared to 25-gauge WH needles. Sharma *et al.*³ and Andres *et al.*⁴ reported that the 26-gauge AT spinal needles were technically easier to use than WH 25-gauge or 27-gauge needles and the PDPH incidences were similar. Sharma further suggested larger studies to better define PDPH and failed spinal blockade rate associated with the AT needle. We, therefore, present here our comparisons of the incidence of PDPH, postdural puncture backache (PDPB) and the failed spinal blockade rate between AT and WH spinal needles.

After Institutional Review Board approval and informed consent, 215 ASA I-II patients undergoing postpartum tubal ligations were randomly assigned to receive subarachnoid anesthesia with either 26-gauge AT (B. Braun Medical, Bethlehem, PA, USA; OD - 0.45 mm, length - 8.89 cm) or 25-gauge WH (Becton-Dickinson, Rutherford, NJ, USA; OD - 0.5 mm, length - 8.89 cm) spinal needles. Spinal needles were introduced with the midline approach at the L2-3, L3-4, or L4-5 intervertebral space. The number of attempts to successful CSF return, success rate of the spinal blockade and the handling characteristics were documented intraoperatively. Postoperatively, an investigator blinded to group assignments interviewed patients daily while in the hospital and then via telephone seven to 14 days after discharge.

The results are summarized in the Table. Incidence of PDPH was similar between AT (3.9%) and WH

TABLE Spinal block characteristics between AT and WH groups (differences between the groups are not significant).

	26-gauge Atraucan AT Group (AT)	25-gauge Whitacre WH Group (WH)
Success with one attempt	62%	61%
Range of number of attempts needed to obtain CSF	1 to 5	1 to 6
Average number of attempts to obtaining CSF	1.5 ± 0.6	1.6 ± 0.7
Patient would accept same spinal technique for future procedures	95%	93%
<i>Quality of spinal blockade</i>		
- Adequate	82%	89%
- Minor supplement	10.6%	4.0%
- Failed	7.7%	7.0%
Unable to identify subarachnoid space	1	2

AT = Atraucan; WH = Whitacre; CSF = cerebral spinal fluid.

(4.0%) groups. The total duration of PDPH was five days and 15 days for AT and WH groups respectively, but the difference in duration between groups did not reach statistical significance. Both groups had a similar one-attempt success rate of 61% (AT) and 62% (WH). The average number of attempts to obtaining CSF were 1.5 ± 0.6 and 1.6 ± 0.7 for AT and WH groups respectively. Incidence of PDPB was 9.6% for AT and 9.0% for WH group; and the severity was mild in both groups. The incidence of failed spinal blockade was 7.7% and 7.0% for AT and WH groups respectively. Ninety-five percent of AT group and 93% of WH group would accept similar spinal anesthesia for future procedures.

In conclusion, we found that the 26-gauge AT and 25-gauge WH spinal needles were associated with similar ease of clinical use and success rate, and a similarly low incidence of PDPH and mild PDPB. The choice between these two types of needles should then be based on the cost and availability of the needles.

Peter H. Pan MD MSEE
Winston-Salem, North Carolina; Richmond, Virginia
Regina Fragneto MD
Lexington, Kentucky
Charlie Moore PhD
Richmond, Virginia
Vernon Ross MD
Winston-Salem, North Carolina
Gina Justis MD
Richmond, Virginia