Correspondence

An additional explanation for atrioventricular block after the administration of atropine

To the Editor:

We wish to comment on the letter by Maruyama *et al.*¹ recently published in the Canadian Journal of Anesthesia. We agree with the pathophysiologic explanation put forward by the authors regarding the imbalance between sympathetic and parasympathetic tone as the cause of the cardiac block observed. We have recently described a series of cases in which nodal rhythm (probably a vagal-mediated anomalous cardiac response similar to sinus arrest) developed in patients with spinal block who were treated with atropine.² The sympathetic blockade that occurs during spinal anesthesia is not the same than during general anesthesia, but a common explanation is possible. Vasodilatation is a general response during both types of anesthesia. In some instances the heart beats without sufficient preload and can develop a paradoxical, predominantly vagal, response consisting of bradycardia or other vagal responses (such as sinus arrest, nodal rhythm, or cardiac arrest).³ In this context, atropine may worsen or potentiate the cardiac response. If our perception is correct, then a sympathomimetic drug (such as epinephrine, as was the choice of the authors, or low-dose ephedrine, the drug used to treat our patients) would be the drug of choice in these cases to increase preload by producing vasoconstriction. In addition ephedrine increases heart rate. This approach has been supported by several authors.^{2,4} Atropine can be a drug of second choice, or perhaps should be administered together with a sympathomimetic drug.

Carlos L. Errando MD PhD
Celsa M. Peiró MD PhD
Valencia, Spain

References

- 1 Maruyama K, Mochizuki N, Hara K. High-degree atrioventricular block after the administration of atropine for sinus arrest during anesthesia (Letter). *Can J Anesth* 2003; 50: 528–9.
- 2 Errando CL. Nodal rhythm after administration of atropine to bradycardic patients under subarachnoid anesthesia. Four cases and a review of pathophysiology and treatment (Spanish). *Rev Esp Anesthesiol Reanim* 2001; 48: 384–6.
- 3 Pollard JB. Cardiac arrest during spinal anesthesia: common mechanisms and strategies for prevention. *Anesth Analg* 2001; 92: 252–6.
- 4 Auroy Y, Bague L, Benhamou D, *et al.* Recommendation of the SOS ALR Group on the use of locoregional anesthesia (French). *Ann Fr Anesth Réanim* 2000; 19: 621–3.

REPLY:

We thank Dr. Errando and Dr. Peiró for their interest in our case report.¹ We reported an abrupt sinus arrest of six seconds duration in a patient under general anesthesia. Atropine 0.5 mg was ineffective in resolving the sick sinus syndrome, and the arrhythmia progressed to atrial flutter with high-degree atrioventricular block.

The fundamental therapeutic challenge of profound bradycardia, as observed in the case reported, or cardiac arrest is successful resuscitation. Vasopressor agents such as ephedrine or epinephrine help maintain preload by their vasoconstrictive action. We attributed the reversal of circulatory collapse to epinephrine's ability to maintain adequate coronary perfusion.

However, we are not sure whether atropine should be considered the second choice of treatment for an arrhythmia caused by an imbalance between sympathetic and parasympathetic tone, even in non-critical cases. The prophylactic administration of vasopressor agents such as ephedrine or epinephrine for such arrhythmias may result in unnecessary hypertension, tachycardia, or ventricular irritability when inhalational anesthesia is administered.² We believe that, similar to treatment of bradycardia during spinal anesthesia,³ the stepwise escalation of treatment of bradycardia from atropine to ephedrine or epinephrine should also be recommended for bradycardia occurring under general anesthesia. We encourage the administration of vasopressor agents together with atropine, especially in life-threatening situations. However, we still feel that the administration of an adequate dose of atropine should be the primary treatment for this type of arrhythmia.

Koichi Maruyama PhD
Noriaki Mochizuki MD
Katsumi Hara MD
Nagano, Japan

References

- 1 Maruyama K, Mochizuki N, Hara K. High-degree atrioventricular block after the administration of atropine for sinus arrest during anesthesia (Letter). *Can J Anesth* 2003; 50: 528–9.
- 2 Pollard JB. Cardiac arrest during spinal anesthesia: common mechanisms and strategies for prevention. *Anesth Analg* 2001; 92: 252–6.
- 3 Johnston RR, Eger EI II, Wilson C. A comparative interaction of epinephrine with enflurane, isoflurane, and halothane in man. *Anesth Analg* 1976; 55: 709–12.

Successful aspiration of blood does not exclude malposition of a large-bore central venous catheter

To the Editor:

We report the occurrence of a primarily unnoticed malpositioned central venous catheter (CVC), in which aspiration of blood failed to indicate CVC tip position in the pleural cavity.

A 44-yr-old man was anesthetized for thoracoscopy-assisted thoracic spinal fusion surgery. After orotracheal intubation the left subclavian vein was punctured, confirmed by the aspiration of abundant dark venous blood. The guide wire and the dilatator passed without resistance and a large-bore multi-lumen catheter (12 Fr $\times$ 8", 3-lumen; Arrow International™, Reading, PA, USA) was subsequently inserted.¹ Aspiration of blood into the transparent part of both large-bore lumina and easy flushing indicated correct intravascular catheter placement. Intraoperatively, when rechecking the catheter, aspiration of blood failed, whereas flushing was easy. Thoracoscopic view revealed a left-side, minimally bloody hydrothorax with the catheter tip freely visible in the intrapleural space (Figure). At the end of surgery the hydrothorax was drained and the misplaced catheter was removed uneventfully.

In the present case, infusion into the pleural cavity was caused by perforation of the subclavian vein. A secondary perforation would explain the successful initial aspiration of blood through both large-bore lines. Yet, most secondary perforations are late complications² and, in addition, erosion is rendered unlikely because practically no blood was detectable in the pleural cavity on thoracoscopy.

Therefore, primary malposition with initial intrapleural placement seems to be more likely.³ Perforation of the subclavian vein with the needle is

FIGURE Thoracoscopic view of the tip of the central venous catheter misplaced in the left pleural cavity.

hardly conceivable, because initial aspiration of blood was abundant. Perforation with the J-shaped guide wire must be considered, as the wire tip can be very stiff and sharp when the tip of the needle is located close to the intima. The clamps on the 12 Gauge (GA) lines do not provide an absolutely tight seal, so blood can fill the distal part of the 12 GA lumina when passing the catheter through the vein lumen. Approximately 0.1 mL blood is enough to produce a continuous 1 cm blood column on aspiration in the proximal transparent part of the catheter. Therefore, it is likely that aspiration of an insufficient blood volume, as evidenced by a blood column in the transparent part of the 12 GA lumina, led to erroneously conclude the catheter was correctly placed. Consequently, the volume of blood aspirated must exceed the volume of the CVC lumina, and an additional control aspiration is recommended immediately before starting the infusion.

Matthias Hohlrieder MD
 Heinrich M. Schubert MD
 Matthias Biebl MD
 Christian Kolbitsch MD
 Patrizia L. Moser MD
 Ingo H. Lorenz MD
 Innsbruck, Austria

References

- 1 Conahan TJ 3rd, Schwartz AJ, Geer RT. Percutaneous catheter introduction: the Seldinger technique (Letter). *JAMA* 1977; 237: 446–7.