vacaine, 10 µg of fentanyl, and 0.2 mg of preservative free morphine at the L2–3 interspace, which was induced with a 25-gauge Pencan needle with the patient in the sitting position. Immediately after the assumption of supine-left-uterine-tilt position an “asymmetric” (a T4 and a T2, on the left and right side, respectively) sensory level of spinal anesthesia with a “window” of unaltered sensation in the mid abdomen, as determined by pinprick, and incomplete motor block of the lower extremities, was reported by the anesthesia resident. Re-evaluation of these findings by the attending anesthesiologist revealed a T10 sensory block (at that time), and upon further questioning the patient admitted to having undergone “cosmetic” breast surgery. Several minutes later the spinal anesthesia reached a T4 sensory level (posterior axillary line) and uneventful abdominal delivery followed. In conclusion, the presumed “asymmetric fast-onset high sensory level” of spinal anesthesia was attributed to the post-mammoplasty breast skin hypoesthesia.

Krzysztof M. Kuczkowski MD
San Diego, California

References

- 1 Ferreira MC, Costa MP, Cunha MS, Sakae E, Fels KW. Sensibility of the breast after reduction mammoplasty. *Ann Plast Surg* 2003; 51: 1–5.
- 2 Reisner LS, Lin D. Anesthesia for cesarean section. In: Chestnut DH (Ed.). *Obstetric Anesthesia: Principles and Practice*. Mosby: St. Louis; 1999: 465–92.

PDA's: are we there yet?

To the Editor:

VanDenKerkhof *et al.*¹ present a fascinating study into the many benefits of personal digital assistants (PDAs) in the clinical setting. I was involved in setting up a PC-based database for anesthetic interventions in a UK maternity hospital. Electronic data capture proved to be much more effective than the use of paper. However, I encountered many problems with the clinical use of PDAs and I would be interested to know of the authors' approach to these.

1. The synchronization process with the PC does not always go smoothly. It is noteworthy that ten sets of study data were lost through synchronization problems.

2. The touchscreens of modern PDAs are very fragile, a colleague cracked the screen of his PDA on the clip of his pager, rendering the PDA useless. It would be naïve to assume the same level of care and attention being shown to a departmental PDA as to a personal one.

3. Since the trial was separate from the acute pain management service round, when brought into full service, how is data transferred to the patient's notes for the patient's care team to review at the ward level?

I have recently worked on a maternity unit where they attempted to use a PDA for data collection, but this was discontinued due to problems one and two outlined above.

Whilst many anesthesiologists in the UK have been keen advocates of PDAs for personal use, I suspect that the technology may not yet be advanced or robust enough for institutional use.

Andrzej Con FRCA
London, UK

Reference

- 1 VanDenKerkhof EG, Goldstein DH, Lane J, Rimmer MJ, Van Dijk JP. Using a personal digital assistant enhances gathering of patient data on an acute pain management service: a pilot study. *Can J Anesth* 2003; 50: 368–75.

REPLY:

We thank Dr. Con for his comments on our article.¹ We respond point by point:

1. We agree that synchronization problems can arise during the downloading process, however, aside from the initial problem in our pilot study, which was related to the structure of the database rather than the synchronization process itself, we encountered no synchronization problems. In a subsequent study, out of 45 assessments we encountered one instance of the personal digital assistant (PDA) freezing during an assessment,² and in another study, out of 360 patient-completed questionnaires (60 on each, a PDA, a Tablet and a Kiosk, and 180 on paper) we encountered two synchronization failures with the PDA and no problems on either the Tablet or the Kiosk.³

2. We encountered no difficulties with the PDA screens during the pilot study; however, the PDAs have been dropped on several occasions over the past three years and the touch screen cracked on one occasion.

3. In our subsequent study where patient assessments were captured on a PDA during acute pain management service rounds, the assessments and orders were sent to a printer using infrared technology.² Printed labels, which had been approved by both patient records, and pharmacy and therapeutics, were affixed to the patient's chart in place of written notes and orders.

Subsequent to the above-mentioned studies, a wireless infrastructure was implemented in our institution.⁴ Wireless technology appears to provide improved security of patient data as no data is stored locally, improved

speed of software running on the PDA and it negates the need for synchronization. Future studies will assess the utility of wireless technology and portable computers at the point-of-care.

Elizabeth VanDenKerkhof DRPH
David H. Goldstein MB BCH MSc FRCPC
Jeremy Lane MD
Michael Rimmer
Janice Van Dijk RN MHSc
Kingston, Ontario

References

- 1 VanDenKerkhof EG, Goldstein DH, Lane J, Rimmer MJ, Van Dijk JP. Using a personal digital assistant enhances gathering of patient data on an acute pain management service: a pilot study. *Can J Anesth* 2003; 50: 368–75.
- 2 VanDenKerkhof EG, Goldstein DH, Rimmer M, Tod D, Kwan Lee H. Handhelds versus paper for acute pain assessments: time and content. *Can J Anesth* 2002; 49: A21 (abstract).
- 3 VanDenKerkhof EG, Goldstein DH, Blaine WC, Rimmer M. Validation of an electronic vs a paper version of the self-completed pre-admission adult anesthetic questionnaire. *Anesth Analg* 2003; 96: S-3 (abstract).
- 4 Goldstein DH, VanDenKerkhof EG, Rimmer MJ. A model for real time information at the patient's side using portable computers on an acute pain service. *Can J Anesth* 2002; 49: 749–54.

Epidural anesthesia in a patient with hyperkalemic periodic paralysis undergoing orthopedic surgery

To the Editor:

Hyperkalemic periodic paralysis (HPP) is a familial abnormality in membrane electrolyte conductance leading to episodes of flaccid weakness in context with shifts in plasma potassium levels. During the attacks plasma potassium is elevated to upper normal levels or above ($> 4.5 \text{ mEq}\cdot\text{L}^{-1}$). Some patients show slight signs of myotonia between and at the beginning of attacks, others display paramyotonia, while in a third category myotonic signs are absent.

A 52-yr-old man with a medical history significant for familial HPP was admitted to the hospital for right hip core decompression. His symptoms of HPP were characterized by two to three attacks/week, lasting one to two hours, and involved his trunk and lower extremities.

On admission, the patient experienced an attack of HPP with muscle weakness and light respiratory distress. Plasma potassium level at this time was $3.4 \text{ mEq}\cdot\text{L}^{-1}$, the electrocardiogram was normal. The patient was hydrated with 5% glucose in water. His SpO_2 on $3 \text{ L}\cdot\text{min}^{-1}$ of oxygen was stable at 98%. Blood pressure, heart rate and rhythm remained stable. The plasma potassium level determined throughout and after surgery ranged between 3.3 to $3.8 \text{ mEq}\cdot\text{L}^{-1}$.

For his surgery, the patient received an epidural anesthetic with 8 mL of lidocaine 2% with epinephrine and 5 mL of levobupivacaine 0.5%. Blood pressure measurements, heart rate and rhythm were stable throughout the procedure. The patient had an overall normal postoperative recovery after epidural anesthesia and his muscle strength returned to baseline during his postanesthesia care unit stay. Until his discharge from hospital, no further episode of HPP occurred.

Overnight fasting, a precipitating factor of HPP, might have contributed to the muscle weakness this patient presented preoperatively. Besides hunger, rest after exercise, cold, infection, and anesthesia have been reported to elicit episodes of HPP. A first report in 1959¹ showed that members of three families developed transient paralysis of two to five hours duration after general anesthesia. Recently, a patient with HPP successfully received hyperbaric bupivacaine 0.75% for spinal anesthesia.² Similarly, the epidural anesthetic with a lidocaine/levobupivacaine mixture in the present patient provided good conditions perioperatively.

In the case here presented, epidural anesthesia was an appropriate anesthetic technique. It provided stable plasma potassium both during surgery and in the immediate postoperative period.

Rima Aouad MD
Peter G. Atanassoff MD
New Haven, Connecticut

References

- 1 Egan TJ, Klein R. Hyperkalemic familial periodic paralysis. *Pediatrics* 1959; 24: 761–73.
- 2 Weller JF, Elliott RA, Pronovost PJ. Spinal anesthesia for a patient with familial hyperkalemic periodic paralysis. *Anesthesiology* 2002; 97: 259–60.