tions, leads one to ask if relatively high concentrations of local anaesthetic should ever be used for postoperative analgesia. Opioids and local anaesthetics have been shown to be synergistic in animal models and at least additive in several human postoperative analgesia trials.²⁻⁴ However, recent reports have shown that an additive analgesic interaction can be attained with relatively low concentrations of bupivacaine (0.1–0.3 mg·ml⁻¹) infused at a rate of 1–5 mg·hr⁻¹.⁽³⁻⁵⁾ It may also be necessary to titrate the local anaesthetic and opioid separately to obtain the greatest degree of analgesic synergy. Fixed mixtures of opioids and local anaesthetics make it difficult to optimize the benefits of each analgesic and reduce side effects.³⁻⁵ This may be best accomplished by utilizing a patient controlled analgesia delivery system for the opioid component.

Kahn's report points out the potential for high concentrations of bupivacaine to produce confusion in the evaluation of postoperative neurological deficits. Concentrations of bupivacaine > 0.06% do not appear to add much in analgesic benefit and contribute to a greater incidence of motor block and hypotension.³⁻⁸ I would suggest that any patient developing motor deficit in the postoperative period should have the local anaesthetic discontinued from the epidural analgesic solution. The patient should be reevaluated within two hours to examine for regression of the nerve block. Furthermore, concentrations of bupivacaine > 0.06% are not warranted and should be abandoned in the delivery postoperative epidural analgesia.

Frederick W. Burgess MD PhD
Providence RI USA

REFERENCES

- 1 Kahn L. Neuropathies masquerading and an epidural complication. *Can J Anaesth* 1997; 44: 313–6.
- 2 Akerman B, Arwestrom E, Post C. Local anesthetics potentiate spinal morphine antinociception. *Anesth Analg* 1988; 67: 943–8.
- 3 Burgess FW, Anderson DM, Colonna D, Cavanaugh DG. Thoracic epidural analgesia with bupivacaine and fentanyl for postoperative thoracotomy pain. *J Cardiothorac Vasc Anesth* 1994; 8: 420–4.
- 4 Burgess FW, Pylman, ML, Helman JD. The ideal epidural bupivacaine concentration for postoperative analgesia. *Anesthesiology* 1993; 79: A796.
- 5 Liu S, Angel JM, Owens BD, Carpenter RL, Isabel, L. Effects of epidural bupivacaine after thoracotomy. *Reg Anesth* 1995; 20: 303–10.
- 6 Badner NH, Komar WE. Bupivacaine 0.1% does not improve postoperative epidural fentanyl analgesia after abdominal or thoracic surgery. *Can J Anaesth* 1992; 39: 330–6.
- 7 George KA, Wright PMC, Chisakuta A. Continuous thoracic epidural fentanyl for post-thoracotomy pain relief: with or without bupivacaine. *Anaesthesia* 1991; 46: 732–6.
- 8 Badner NH, Reimer EJ, Komar WE, Moote CA. Low-dose bupivacaine does not improve postoperative epidural fentanyl analgesia in orthopedic patients. *Anesth Analg* 1991; 72: 337–41.

REPLY

Synergistic effects of combined opioid-local anaesthetic epidural mixtures are more evident in thoracic than in lumbar epidural¹ infusions. However, the optimum concentration of each drug has yet to be determined. In an ideal setting, titration of each drug could best achieve the desired result. This may be impractical on a busy Acute Pain Service, and expensive, as two infusers and two drug mixtures are required. Our experience with very low concentrations (0.01–0.03%) of bupivacaine resulted in inconsistent analgesia, depending on the surgery, the site of the epidural catheter, drug mixtures and infusions rates. In addition it is labour intensive. Despite these low concentrations, sensorimotor blocks with lumbar epidural catheters still occur.²

The references quoted to reduce concentrations of bupivacaine do not support the recommendations. Reference #3, by addressed the fentanyl sparing effects of various concentrations of bupivacaine with no mention of the deleterious effects of bupivacaine 0.125%. Although lower concentrations of bupivacaine were used, high infusion rates of up to 22 ml·hr⁻¹ were needed, raising the question, was less drug used? Reference #4³ is an abstract, non-peer reviewed article, with few patients enrolled in each group. Reference #7⁴ et al. suggested using bupivacaine 0.2% with fentanyl in thoracic epidurals for thoracotomies. References #6⁵ and #8⁶ demonstrated that low dose bupivacaine did not improve analgesia. Perhaps higher doses are needed. In a follow up study,⁷ bupivacaine 0.125% was found to improve analgesia after thoracic and abdominal surgery, with minimal sensorimotor disturbance.

Using 8–12.5 mg·hr⁻¹ epidural bupivacaine as a continuous infusion would not be considered high dose. In a recent review Stevens and de Leon-Casasola,⁸ recommended 6–10 mg·hr⁻¹ bupivacaine 0.2% for epidural infusions for postoperative analgesia. When used according to the guidelines in the case report,⁹ the incidence of motor blocks was low. We have found this regimen to pro-

vide analgesia reliably and safely to the majority of our postoperative patients.

To recommend that "concentrations of bupivacaine > 0.06% are not warranted and should be abandoned in the delivery of postoperative epidural analgesia" is not based on sound evidence. Perhaps ropivacaine,¹⁰ with its motor sparing qualities, may offer some solution. Vigilance, frequent monitoring, titrated rather than constant infusions¹¹ and appropriate management of motor block should be an integral part of continuous epidural infusion care.

Larry Kahn MB CHB
Hamilton, Ontario

REFERENCES

- 1 Burgess FW, Anderson DM, Colonna D, Cavanaugh DG. Thoracic epidural analgesia with bupivacaine and fentanyl for postoperative thoracotomy pain. *J Cardiothorac Vasc Anesth* 1994; 8: 420-4.
- 2 Youngsrom P. Dilute local anesthetics in epidural analgesic solution (Letter). *Anesth. Analg.* 1992; 74: 313.
- 3 Burgess FW, Pylman ML, Helman JD. The ideal epidural bupivacaine concentration for postoperative analgesia. *Anesthesiology* 1993; 79: A796.
- 4 George KA, Wright PMC, Chisakuta A. Continuous thoracic epidural fentanyl for post-thoracotomy pain relief: with or without bupivacaine. *Anaesthesia* 1991; 46: 732-6.
- 5 Badner N, Komar W. Bupivacaine 0.1% does not improve post-operative epidural fentanyl analgesia after abdominal or thoracic surgery. *Can J Anaesth* 1992; 39: 330-6.
- 6 Badner NH, Reimer EJ, Komar WE, Moote CA. Low-dose bupivacaine does not improve postoperative fentanyl analgesia in orthopedic patients. *Anesth Analg* 1991; 72: 337-41.
- 7 Badner NH, Bhandari R, Komar WE. Bupivacaine 0.125% improves continuous postoperative epidural fentanyl analgesia after abdominal or thoracic surgery. *Can J Anaesth* 1994; 41: 387-92.
- 8 Stevens RA, de Leon-Casasola O. What have we learned about acute postoperative epidural pain management since 1988? *Techniques in Regional Anesthesia and Pain Management* 1997; 1: 59-63.
- 9 Kahn L. Neuropathies masquerading as an epidural complication. *Can J Anaesth* 1997; 44: 313-6.
- 10 Scott DA, Chamley DM, Mooney PH, Deam RK, Mark AH, Hagglof B. Epidural ropivacaine infusion for postoperative analgesia after major lower abdominal surgery—a dose finding study. *Anesth Analg* 1995; 81: 982-6.
- 11 Kahn L, Jackson P. Fentanyl infusions for post-thoracotomy pain management (Letter). *Can J Anaesth* 1994; 41: 871.

Detecting endobronchial intubation during breast surgery

To the Editor:

A 69-yr-old lady (weight 79 kg, height 148 cm) was anaesthetised for breast reduction surgery. After induction, intubation, and the institution of mechanical ventilation, vital signs, recorded on a Datex AS3 monitor, were stable (see Table below). Breath sounds were normal and equal, so the low oxygen saturation was accepted, given her age, weight and smoking history. After 12 min, well into the dissection of the left breast, a decrease in oxygen saturation, increase in airway pressure, and decrease in compliance, all pointed to endobronchial intubation.

Before retracting the endotracheal tube, I wished to confirm the diagnosis by auscultation. However, all of the front and sides of the patient's chest were prepped and draped, and part of the surgical field.

Previously, the surgeon had discussed this as a hypothetical situation and the solution suggested was to place a stethoscope in a sterile bag. A clear plastic instrument pouch was provided by the scrub nurse, who positioned the stethoscope in each axilla. No breath sounds were heard on the left side, confirming the diagnosis. The endotracheal tube was pulled back 2 cm, breath sounds returned, and the vital signs returned to their previous values.

Acknowledgments: To Drs John Ascah and John Toye, and Theresa Tobson RN

John Oyston MBBS FFARCS
Orillia Soldiers' Memorial Hospital
Orillia, Ontario

TABLE

Time	SpO ₂ %	P _{ET} CO ₂ mmHg	Compliance ml·cmH ₂ O ⁻¹	PAirway cm H ₂ O	Heart bpm	BP mmHg
08:25	93	46	32	23	81	128/80
08:37	86	45	21	26	75	181/109
08:42	93	46	32	23	99	168/94