William P.S. McKay MD FRCP
Saskatoon, Saskatchewan

References

- 1 Saitoh Y, Nakajima H, Hattori H, Aoki K, Katayama T, Murakawa M. Neuromuscular blockade can be assessed accelerographically over the vastus medialis muscle in patients positioned prone. *Can J Anesth* 2003; 50: 342–7.
- 2 Warwick R, Williams PL. *Gray's Anatomy*, 35th ed. Edinburgh: Longman; 1973: 565.
- 3 Basmajian JV. *Grant's Method of Anatomy*, 10th ed. Baltimore: Williams & Wilkins; 1980: 254.

REPLY:

We studied the accelerographic monitoring of neuromuscular block over the vastus medialis muscle.¹ However, we were unaware of the precise innervation of the vastus medialis muscle. As suggested, in our study, small im branches of the vastus medialis nerve may have been stimulated. Further studies will be required to evaluate the appropriateness of direct muscle stimulation to monitor neuromuscular blockade in anesthetized patients. The degree of neuromuscular block assessed accelerographically over the directly stimulated muscle should be compared to that evaluated mechanically, electromyographically, or accelerographically at the thumb or the adductor pollicis muscle. We thank Dr. McKay for his constructive suggestion.

Yuhji Saitoh MD
Fukushima, Japan

Reference

- 1 Saitoh Y, Nakajima H, Hattori H, Aoki K, Katayama T, Murakawa M. Neuromuscular blockade can be assessed accelerographically over the vastus medialis muscle in patients positioned prone. *Can J Anesth* 2003; 50: 342–7.