Frances Chung, University of Toronto, University Health Network, Toronto, ON, Canada;

Balaji Yegneswaran, University of Toronto, University Health Network;

F Herrera, University of Toronto, University Health Network;

A Shenderey, University of Toronto, University Health Network;

C Shapiro, University of Toronto, University Health Network;

Background: Obstructive sleep apnea (OSA) and difficult tracheal intubations are two significant clinical problems that contribute to increased morbidity and mortality among surgical patients. In a small retrospective study of 15 patients, Hiremath et al found the prevalence of OSA was 53% in patients with difficult intubation(1). We hypothesized that patients who presented with difficult intubation might have a high incidence of OSA and patients with difficult intubation should be referred for Polysomnography (PSG) studies.

Methods: Hospital ethics approval was obtained. Patients classified as grade 3 and 4 according to Cormack and Lehane direct laryngoscopic view or patients who required two or more attempts for successful endotracheal intubation were referred by anesthesiologists at four hospitals.

Informed consent for overnight polysomnography was obtained from these patients when they had recovered from surgery. The polysomnography was conducted 6-8 weeks after surgery to avoid the effect of postoperative recovery. Data on Apnea-Hypopnea Index (AHI), oxygen saturation and postoperative complications were collected. American Academy of Sleep Medicine defines mild OSA: AHI 5-15/hr, moderate OSA: AHI 15-30/hr, severe OSA: AHI>30/hr. Clinical and polysomnographic variables were compared between patients with difficult intubation with and without OSA using t-tests.

Results: Over a twenty-month period, 84 patients with difficult intubations were referred by anesthesiologists. Thirty three patients agreed to participate. The mean age of patients who underwent polysomnography was 57±13 yrs (range: 22-78 years), ASA I-9.1%; II-36.4%; III-50.5%; IV-3%. 66%(22/33) of the patients with difficult intubation had an AHI>5/hr. 30%(10/33) were mild OSA, 18%(6/33) moderate OSA and 18%(6/33) severe OSA. There were significant differences ($p<0.05$) in gender, neck size and quality of sleep between the OSA group and non-OSA group. The overall incidence of postoperative events was 21%(7/33). Of the 7 complications 86%(6/7) complications were among OSA patients. 18%(4/22) of OSA patients had severe desaturation during their postoperative period and 32%(7/22) of OSA patient needed oxygen therapy during their stay in the ward. There was no unplanned ICU admission among the patients studied.

Discussion: 66% of patients with difficult intubation were diagnosed to have OSA by polysomnography. Since untreated OSA patients have a shortened lifespan by twenty years(2), anesthesiologists have an obligation to refer patients with difficult intubation for PSG.

Reference:

1. BJA 1998 80: 606-611
2. Thorax 1998 Suppl 3: S16-9

Table – Demographic and polysomnographic data

	AHI>5/hr (n- 22)	AHI ≤5/hr (n – 11)
Age (year)	60±10	50±17
Gender (Male/Female)	18/4*	5/6
BMI (kg/m²)	32±7	28±8
Neck Size (cm)	42±4*	37±4
AHI-TST	26 ± 22*	2±1
AHI-REM	33±24*	10±10
Arousal Index	32±15*	16±9
Min. SaO₂	79.±10*	90±6

Mean±SD * p<0.05

AHI-TST – Apnea Hyponea Index Total Sleep Time

AHI-REM - Apnea Hyponea Index Rapid Eye Movement Sleep