model suffers from over-fitting: internal validation such as bootstrapping, and external validation on a new group of patients.^{4,5} The authors did not validate their model. Their statement that the model's high ROC area implies reproducibility is false: in this setting, ROC area simply measures the accuracy of the model when it is reapplied to the sample population; it has no relation to reproducibility or external validity.⁶

Keyvan Karkouti MD FRCPC MSC
Keith Rose MD FRCPC
Marsha Cohen MD PhD
Dan Wigglesworth BSc
Toronto, Canada

References

- 1 Naguib M, Malabarey T, AlSatli RA, Damegh SA, Samarkandi AH. Predictive models for difficult laryngoscopy and intubation. A clinical, radiologic and three-dimensional computer imaging study. *Can J Anesth* 1999; 46: 748–59.
- 2 Bellhouse CP, Doré C. Criteria of estimating likelihood of difficulty of endotracheal intubation with the Macintosh laryngoscope. *Anaesth Intensive Care* 1988; 16: 329–37.
- 3 Harrell FE Jr, Lee KL, Mark DB. Multivariable prognostic models: issues in developing models, evaluating assumptions and adequacy, and measuring and reducing error. *Stat Med* 1996; 15: 361–87.
- 4 Wasson JH, Sox HC, Neff RK, Goldman L. Clinical prediction rules. Applications and methodological standards. *N Engl J Med* 1985; 313: 793–9.
- 5 Laupacis A, Sekar N, Stiell IG. Clinical prediction rules. A review and suggested modifications of methodological standards. *JAMA* 1997; 277: 488–94.
- 6 Feinstein AR. Multiple logistic regression. In: Feinstein AR (Ed.). *Multivariable Analysis: An Introduction*. New Haven: Yale University Press, 1996: 297–330.

REPLY:

We appreciate Dr. Karkouti's interest in our study.¹ Dr Karkouti did not provide any evidence to support his contention that "our clinical model may not be applicable for use in clinical practice."

We suspect that the head and neck movement (as part of the Wilson risk sum score) and the radiological measurements (with the head erect and mouth closed, and with the head fully extended on the neck and the mouth fully) represent atlanto-occipital extension. We addressed in our paper the reliability of the Wilson risk sum score. We included all of the radiological measurements described by Bellhouse and Doré.² Therefore, we do not follow Dr. Karkouti's statement regarding Bellhouse's study.

Of importance, our model was validated with bootstrapping. Our model was developed on data from 65 patients and not only from the 25 patients who were difficult to intubate. Dr. Karkouti's comments on the ROC curve are incorrect. For detailed discussion on the area under ROC curve, see Hanley and McNeil.³ We are currently conducting a clinical study on a different population of patients to evaluate whether the scale applies.

Mohamed Naguib MD
Iowa City, Iowa, USA

References

- 1 Naguib M, Malabarey T, AlSatli RA, Al Damegh S, Samarkandi AH. Predictive models for difficult laryngoscopy and intubation. A clinical, radiologic and three-dimensional computer imaging study. *Can J Anesth* 1999; 46: 748–59.
- 2 Bellhouse CP, Doré C. Criteria for estimating likelihood of difficulty of endotracheal intubation with the Macintosh laryngoscope. *Anaesth Intensive Care* 1988; 61: 329–37.
- 3 Hanley JA, McNeil BJ. The meaning and use of the area under a receiver operating characteristic (ROC) curve. *Radiology* 1982; 143: 29–36.

Is aspiration prophylaxis necessary before spinal anesthesia for Cesarean section. ?

To the Editor:

Failure of spinal anesthesia (SA) for Cesarean section (CS) necessitating general anesthesia (GA) and tracheal intubation is extremely low. However, aspiration prophylaxis (AP) is recommended because of the possibility of either failure or excessive spread of spinal block requiring GA. We have reviewed, retrospectively, the penoperative course of parturients undergoing CS under SA at our hospital from 1991 to 1993. When spinal block failure occurred, sodium bicarbonate *po* was administered prior to GA. Among the 743 cases that we reviewed, failure occurred in 15 cases (2%) because of inadequate analgesia (1.9%) or prolonged surgery for hysterectomy (0.1%). No case required intubation due to excessive block. No pulmonary aspiration was identified. In a survey of 165 academic departments in the USA and Canada, Greenhalgh¹ found that 94% of elective and 99% of emergency CS patients received aspiration prophylaxis. Plumer² found that 88% of the anesthesiologists attending a major obstetric anesthesia conference in the USA routinely gave prophylaxis