Correspondence

Cannabinoids for postoperative analgesia?

To the Editor:

We read with interest the article on the effects of nabilone on postoperative pain by Pierre Beaulieu.¹ While it is unfortunate that his results did not generate firm conclusions, we believe that descriptive subgroup analyses may have yielded some important information regarding the effects of nabilone on postoperative morphine consumption and side-effects. Nineteen of the 41 patients in the study underwent hysterectomy or myomectomy, which is a surgically homogenous patient population for whom it would have been interesting to have compared the effects of placebo, ketoprofen and nabilone doses of 1 and 2 mg on morphine consumption, visual analogue scale (VAS) scores and secondary outcomes. Eighteen of the 41 patients underwent orthopedic surgery (hip or knee arthroplasty), which represents a different surgical population of potentially greater heterogeneity, with an additional confounding factor that knee arthroplasties are associated with more postoperative pain compared with hip arthroplasty.² A greater number of knee arthroplasty patients in the nabilone 2 mg group could explain why there is an apparent possible hyperalgesic effect of nabilone. Hence, it would be interesting to know the breakdown of knee/hip arthroplasty by study group (placebo, ketoprofen and nabilone 1 and 2 mg), and the results (morphine consumption, VAS scores and secondary outcomes) according to the orthopedic procedure and study group.

Étienne de Médecis MD MSc FRCPC

Jean-Pierre Tétrault MD MSc FRCPC

Centre hospitalier universitaire de Sherbrooke,
Sherbrooke, Canada

E-mail: estrial@globetrotter.net

Accepted for publication November 6, 2006.

References

- 1 Beaulieu P. Effects of nabilone, a synthetic cannabinoid, on postoperative pain. *Can J Anesth* 2006; 53: 769–75.
- 2 Pang WW, Hsu TC, Tung CC, Hung CP, Chang DP, Huang MH. Is total knee replacement more painful than total hip replacement? *Acta Anaesthesiol Sin* 2000; 38: 143–8.

Reply:

Thank you to Dr. de Médecis and Dr. Tétrault for their interesting comments. While it is true that recruited patients in the referenced study underwent primarily gynecological and orthopedic procedures, the patients were distributed into four different groups, making it difficult to perform appropriate sub-group analysis with the limited number of subjects in each group.

The suggestion that more patients in the nabilone 2-mg group underwent knee arthroplasty is in fact, correct. Nevertheless, if excluding the three patients who underwent knee arthroplasty, the amount of morphine patient-controlled analgesia used in 24 hr (the primary outcome variable) in the nabilone 2-mg group was higher than previously reported (average of 49.5 mg compared to 45.4 mg, with no significant differences between the four groups). Furthermore, the intensity of pain at rest (24 hr), which was significantly different between groups, remained so, even if knee arthroplasty patients were excluded from the analysis. In contrast, when considering the pain scores on movement (24 hr), there were no statistical differences, although mean pain scores in the nabilone 2-mg group remained numerically greater, and only decreased from 7.7 to 7.1 ($P = 0.09$). Therefore, despite the fact that more patients in the nabilone 2-mg group underwent knee arthroplasty, this fact did not markedly influence pain scores which remained highest in this group, despite the similarity of morphine consumption in the two groups.

Pierre Beaulieu MD PhD

CHUM – Hôtel-Dieu, Montreal, Canada

E-mail: pierre.beaulieu@umontreal.ca

Angioedema, angiotensin converting enzyme inhibitors, and angiotensin receptor blocking drugs

To the Editor:

We read the review by Sarkar *et al.*¹ with great interest. Although angioedema can cause rapid, progressive airway compromise, facial and oral cavity edema as a complication of angiotensin converting enzyme