realize that these are interim guidelines - a stepping stone to a more formal Canadian certification process - but in this one regard we think the recommendations are overly restrictive and ignore what a dedicated, enthusiastic group can do within their current framework of practice.

Craig Bosenberg MB ChB FRCPC

Joy Dobson MD FRCPC

Stan Vuksic MD FRCPC

Regina Qu'Appelle Health Region, Regina, Canada

E-mail: vuksics@sasktel.net

Accepted for publication November 15, 2006.

References

- 1 Beique F, Ali M, Hynes M, et al. Canadian guidelines for training in adult perioperative transesophageal echocardiography. *Can J Anesth* 2006; 53: 1044-60.
- 2 Finegan BA. Progress through cooperation: securing a sound training pathway for perioperative transesophageal echocardiography. *Can J Anesth* 2006; 53: 969-72.

Reply:

We thank Dr. Bosenberg, Dr. Dobson and Dr. Vuksic for their interest in the recently published Canadian guidelines for training in adult perioperative transesophageal echocardiography (TEE).¹ The consensus panel discussed in great detail the importance of dedicated echocardiography training to achieve a basic level of expertise. Several points need to be emphasized to further explain the recommendations outlined in the Canadian guidelines. First, it is important that trainees receive dedicated and concentrated exposure to TEE unencumbered by other clinical responsibilities and distractions. With the current human resource shortage, such distractions can easily overshadow the trainee's needs. Our experience with trainees suggests that intermittent TEE exposure during training with inconsistent TEE attendance resulted in poor grasp of the necessary concepts, poor skills at probe manipulation, and slow performance of the echocardiographic examination. Teachers may also feel frustrated due to inefficient repetitions of previously taught principles that were forgotten when there is only sporadic exposure to TEE. Moreover, as stated in the guidelines, although the exact duration of training is not as important as completing the required number of echocardiographic examinations and continuing medical education (CME) hours, the proposed three months of training at a basic level is consistent with previously published cardiology guidelines for training in echocardiography.²⁻⁴

Achieving a basic level of expertise is rewarded in the Canadian guidelines with a level of autonomy which enables the independent echocardiographic assessment of cardiac pathology and surgical procedures including uncomplicated valve surgery. Consultation with a physician who has achieved an advanced level of expertise should only be necessary in more complex cases or when new TEE findings may result in a modification of the surgical procedure. Therefore, training at a basic level must be adequate for the trainee to have the ability to identify and recognize cardiac findings that may be significant and would require consultation. A wrong diagnosis or failure to identify important cardiac pathology can have catastrophic consequences for the patient. It is essential that this level of autonomy is associated with stringent training criteria as outlined in the Canadian guidelines. Moreover, the issue of consultation when abnormalities are detected should be viewed as a positive learning experience that will improve patient care. This process should also allow physicians with basic training to further their knowledge and potentially progress to an advanced level of expertise in parallel with clinical activity as described in the Canadian guidelines.¹ Even advanced practitioners regularly engage in this exercise when confronted with more complex cardiac cases.

It is not the intent of the panel to dismiss alternate training pathways. In fact, we have stated "The current consensus acknowledges that there are several valid training pathways to achieve each level of expertise in perioperative echocardiography. The Canadian training guidelines in perioperative TEE only highlight some of the possible training pathways that can be used to facilitate this process."¹ The emphasis is on obtaining the appropriate knowledge and skills for each level of training in perioperative TEE and not on the specific pathway that is used to achieve this goal. What is proposed in the Canadian guidelines is a reliable pathway with a concrete number of echocardiographic studies, continuing medical education (CME) hours and duration of training to produce a competent echocardiographer. In addition under "Grandfathering of established practitioners" we have recognized that "there are highly qualified practitioners actively engaged in perioperative TEE whose training did not follow the guidelines outlined in this document but who nonetheless have learned the techniques in other ways and have attained a high level of competence. It is not the intention of this document to prevent such individuals from practicing and providing important clinical care. It is hoped that this document will provide hospitals and regulatory committees general guidelines by which such nonstandard training can be evaluated."¹ Expertise in TEE is difficult to achieve and mandates appropriate training to ensure that responsibilities are commensurate with the level of training.

François Béique MD FRCPC*
 Mohamed Ali MD FRCPC
 Scott MacKenzie MD FRCPC
 Mark Hynes MD FRCPC
 SMBD Jewish General Hospital,* Montreal, Canada
 E-mail: fbeique@ana.jgh.mcgill.ca

References

- 1 Béique F, Ali M, Hynes M, et al. Canadian guidelines for training in adult perioperative transesophageal echocardiography. *Can J Anesth* 2006; 53: 1044–60.
- 2 Pearlman AS, Gardin JM, Martin RP, et al. Guidelines for optimal physician training in echocardiography. Recommendations of the American Society of Echocardiography Committee for Physician Training in Echocardiography. *Am J Cardiol* 1987; 60: 158–63.
- 3 Stewart WJ, Aurigemma GP, Bierman FZ, et al. Guidelines for training in adult cardiovascular medicine. Core Cardiology Training Symposium (COCATS). Task Force 4: training in echocardiography. *J Am Coll Cardiol* 1995; 25: 16–9.
- 4 Quinones MA, Douglas PS, Foster E, et al. American College of Cardiology/American Heart Association Clinical Competence Statement on Echocardiography: a report of the American College of Cardiology/American Heart Association/American College of Physicians – American Society of Internal Medicine Task Force on Clinical Competence. *Circulation* 2003; 107: 1068–89.

A simple technique to secure the endotracheal tube over an intubating fibroscope

To the Editor:

We use the Intubation Fiberscope (Pentax FI-10P2; Tokyo, Japan) in patients with anticipated difficult airways and in situations where neck movement should be avoided during intubation, including all causes of atlantoaxial instability. The technique of intubation with the fibroscope consists of placing the shaft of the bronchoscope tube in the trachea under indirect vision, followed by railroading the endotracheal tube over the insertion tube. Before intubation, the endotracheal tube is loaded on the bronchoscope up to the proximal end of the insertion tube of the fibroscope. The endotracheal tube is then secured at the proximal end by sliding up the tube firmly, with its position being maintained by a twisting motion or use of adhesive tape. However, not infrequently, the endotracheal tube slides down the fibroscope shaft during bronchoscopy, and additional assistance may be required to

FIGURE The endotracheal tube is held in position by a rubber band sling, with a figure-of-eight around the blue connector and instrument channel inlet; the inset shows an enlarged view of the blue connector of the endotracheal tube and instrument channel on the control body of the intubation fibroscope.

secure its position during the endoscopic procedure to prevent interference and to facilitate proper positioning of the fibroscope within the trachea.

We have adapted a simple technique that easily and reliably secures the endotracheal tube at the proximal end of the insertion tube. The fibroscope control-body has an instrument channel, the proximal end of which protrudes like a hook. From this instrument channel, the endotracheal tube may be easily secured by placing a rubber band at its proximal end around the 15 mm blue connector. Thereafter, the endotracheal tube is threaded over the shaft of the intubation fibroscope. Once positioned, the rubber band is rotated to achieve a figure-of-eight, and is then slid over the proximal end of the instrument channel (Figure). The rubber band firmly secures the endotracheal tube in place, and the tube can be released easily after the fibroscope has been correctly positioned. We have found this method of temporary securing of the endotracheal tube to be effective and reliable during fiberoptic intubation.