TABLE Experience with GlideScope® for tracheal intubation in neonates

Case	Sex	Birth weight (g)	Gestational age (weeks)	Primary respiratory disease	Indication for tracheal intubation	Tracheal intubation with GlideScope® video laryngoscope
1	M	3500	38	Congenital diaphragmatic hernia	Progressive respiratory failure	Yes
2	M	2970	36	Respiratory distress syndrome	Surfactant administration	Yes
3	F	2260	32	Respiratory distress syndrome	Surfactant administration	No
4	M	590	24	Respiratory distress syndrome	Tracheal tube exchange	No
5	M	590	23	Respiratory distress syndrome	Tracheal tube exchange	Yes

for this group of preterm infants, development of a smaller version, suitable for neonatal use, may be warranted.

The two main advantages of the GVL, when compared to standard laryngoscopes, are an improved view of the larynx facilitating successful tracheal intubation, and a potential role for teaching purposes.³⁻⁵ While these aspects are important for personnel dedicated to adults patients, they could assume equal, if not greater, potential importance for physicians involved in the care of the neonatal airways.

Finally, we noticed that the position of the operator using GVL is different from operator positioning for direct laryngoscopy. The orientation of the videochip imparts a different laryngoscopic view as compared to that seen when simultaneously looking into the mouth.⁴ Our experience suggests the requirement for operator training with this new instrument before routine use. On a positive note, the manipulations of the GVL by the resident, and the observed effect of these manipulations on the position of the blade tip on the GVL display and passage of the TT made it a valuable teaching device. The verbal instructions to the resident were also helpful to attending observers.

In conclusion, we feel that the GVL could become an effective device for neonatal intubation. However, the neonatal model GVL we used was not entirely satisfactory in the neonatal population. This group of patients may well benefit from further refinements of the GVL specifically designed for neonatal use.

Daniele Trevisanuto MD*

Enrica Fornaro MD*

Chandy Verghese MD†

University of Padova,* Azienda Ospedaliera di Padova, Padova, Italy

Royal Berkshire Hospital National Health Service Trust,† Reading, United Kingdom

E-mail: trevo@pediatria.unipd.it

Accepted for publication December 5, 2005.

References

- 1 Sun DA, Warriner CB, Parsons DG, Klein R, Umedaly HS, Moulton M. The GlideScope® Video Laryngoscope: randomized clinical trial in 200 patients. *Br J Anaesth* 2005; 60: 94–381.
- 2 Cuchillo JV, Rodriguez MA. Considerations aimed at facilitating the use of the new GlideScope® videolaryngoscope (Letter). *Can J Anesth* 2005; 52: 661–2.
- 3 Rai MR, Dering A, Verghese C. The GlideScope® system: a clinical assessment of performance. *Anaesthesia* 2005; 60: 60–4.
- 4 Cooper RM. Use of a new videolaryngoscope (GlideScope®) in the management of a difficult airway. *Can J Anesth* 2003; 50: 611–3.
- 5 Doyle DJ. GlideScope®-assisted fiberoptic intubation: a new airway teaching method (Letter). *Anesthesiology* 2004; 101: 1252.

Verifying spinal needle location in the presence of a “dry tap”

To the Editor:

It is with great interest that I read the letter by Ramachandran and Ponnusamy describing successful spinal anesthesia after multiple attempts with a distinct “give”, but no free flow of cerebrospinal fluid (CSF).¹ The authors of this article should be commended for their cautious management of this infrequent clinical situation. I agree with the authors’ statement that “common sense dictates that the procedure be repeated, but if the outcome remains the same, and the patient refuses general anesthesia the options are limited.” I would also like to point out other options that may be considered. Although radiological imaging with contrast may be impractical and cumbersome in some settings, this diagnostic approach would verify exact needle position prior to injection of the local anesthetic. The recently introduced epidural stimu-

lation test can also provide objective assessment of needle location, without cumbersome equipment.

Electrical stimulation is a useful and reliable 'real-time' technique to confirm epidural insulated needle placement.²⁻⁵ Not only do muscle twitches elicited by low current stimulation detail the position of the insulated needle within the epidural space, but the threshold current for motor responses can also be used to predict intrathecal placement.²⁻⁵ Insulated needles correctly situated within the epidural space should result in muscle twitches being elicited with a current well above 1 mA. However, whenever a motor response is detected at currents below or around 1 mA, intrathecal, or subdural placement should be suspected. Alternatively, the needle may be resting against a nerve. Under very low current stimulations, intrathecal needle placement will likely produce a bilateral motor response. Subdural placement may elicit multiple patchy segmental responses; whereas needles placed against a nerve root will elicit a unilateral single level motor response. In addition, paresthesiae are likely to occur if the spinal needle is placed directly against a nerve root in an awake patient. The advantage of real-time stimulation testing is that needle location can be assessed objectively, even in the absence of CSF flow, before injecting a loading dose of local anesthetic.

The clinical response to the epidural stimulation test using an *insulated* needle, in the absence of free flow of CSF, has not been reported. This scenario might reproduce the local motor responses associated with intrathecal placement of an insulated needle at low current (< 1 mA). I would caution that experience with the technique is based upon results using an *insulated* needle or catheter, whereas commonly used spinal needles are *uninsulated*. Thus, even if a confirmation test had been utilized, the authors' assertion that the (spinal anesthesia) procedure be repeated, still holds true.¹

Ban C.H. Tsui MSC MD FRCP(C)
University of Alberta Hospital, Edmonton, Canada
E-mail: btsui@ualberta.ca
Accepted for publication December 17, 2005.

References

- 1 Ramachandran K, Ponnusamy N. Dry tap and spinal anesthesia (Letter). *Can J Anesth* 2005; 52: 1104-5.
- 2 Tsui BC, Tarkkila P, Gupta S, Kearney R. Confirmation of caudal needle placement using nerve stimulation. *Anesthesiology* 1999; 91: 374-8.
- 3 Tsui BC, Wagner A, Cave D, Seal R. Threshold current for an insulated epidural needle in pediatric patients. *Anesth Analg* 2004; 99: 694-6.
- 4 Tsui BC, Wagner AM, Cunningham K, Perry S, Desai S, Seal R. Threshold current of an insulated needle in the intrathecal space in pediatric patients. *Anesth Analg* 2005; 100: 662-5.
- 5 Tsui BC, Wagner AM, Cunningham K, Perry S, Desai S, Seal R. Can continuous low current electrical stimulation distinguish insulated needle position in the epidural and intrathecal spaces in pediatric patients? *Paediatr Anaesth* 2005; 15: 959-63.